

Appendix A: Search Strategies

Number of citations in ()

/ after an index term indicates that all subheadings were selected.

* before an index term indicates that that term was focused - i.e. limited to records where major MeSH/Emtree term.

"exp" before an index term indicates that the term was exploded.

.tw. indicates a search for a term in title/abstract.

.mp. indicates a free text search for a term.

.pt. indicates a search for a publication type.

\$ at the end of a term indicates that this term has been truncated.

? in the middle of a term indicates the use of a wildcard.

adj indicates a search for two terms where they appear adjacent to one another.

sh indicates a search term for subheading.

MEDLINE (OVID) for Randomized Controlled Trials Using the Cochrane Highly Sensitive and Specific Search Strategy (Sensitivity and Precision Maximizing Version 2008)

1. Coronary Artery Disease/ or Coronary Disease/
2. Myocardial Ischemia/
3. Angina Pectoris/ or Angina, Unstable/
4. Angina Pectoris/ or Arterial Occlusive Diseases/
5. Peripheral Vascular Diseases/
6. Vascular Diseases/
7. Atherosclerosis/
8. Cardiovascular Diseases/
9. Carotid Artery Diseases/
10. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
11. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10
12. randomized controlled trial.pt.
13. controlled clinical trial.pt.
14. randomized.ab.
15. placebo.ab.
16. clinical trials as topic.sh.
17. randomly.ab.
18. trial.ti.
19. 12 or 13 or 14 or 15 or 16 or 17 or 18
20. humans.sh.
21. 19 and 20
22. (alacepril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril or moveltipril or pentopril or perindopril or quinapril or ramipril or spirapril or temocapril or teprotide ortrandolapril or zofenopril).mp.
23. (losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan).mp.
24. Angiotensin-Converting Enzyme Inhibitors/
25. Angiotensin II Type 1 Receptor Blockers/
26. (ACEI or ARB).mp.

Appendix A: Search Strategies

27. 22 or 23 or 24 or 25 or 26

28. 11 and 21 and 27

Appendix A: Search Strategies

CENTRAL (OVID) for Randomized Controlled Trials

1. (alacepril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril).mp.
2. (moveltipril or pentopril or perindopril or quinapril or ramipril or spirapril or temocapril or teprotide ortrandolapril or zofenopril).mp.
3. (losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan).mp.
4. Angiotensin-Converting Enzyme Inhibitors/
5. Angiotensin II Type 1 Receptor Blockers/
6. (ACEI or ARB).mp.
7. 1 or 2 or 3 or 4 or 5 or 6
8. Coronary Artery Disease/ or Coronary Disease/
9. Myocardial Ischemia/
10. Angina Pectoris/ or Angina, Unstable/
11. Arterial Occlusive Diseases/
12. Peripheral Vascular Diseases/
13. Vascular Diseases/
14. Atherosclerosis/
15. Cardiovascular Diseases/
16. Carotid Artery Diseases/
17. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
18. 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17
- 19. 7 and 18**

Appendix A: Search Strategies

EMBASE (Silver Platter) for Randomized Controlled Trials Using the McMaster Health Information Research Unit (HiRU) highly sensitive, highly specific EMBASE search strategy for treatment articles that minimizes differences between sensitivity and specificity

((((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)) and (((angiotensin converting enzyme inhibitor or ACE inhibitor or ACEI or arb or angiotensin receptor blocker or angiotensin ii type 1 receptor blocker) or (alacapril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril or moveltipril or pentopril or perinodopril or quinapril or ramipril or spirapril or temocapril or teprotide or trandolapril or zofenopril or losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan)) and (((random) in AB) or (double-blind) or (placebo) or ((random) in TI)))

Appendix A: Search Strategies

MEDLINE (OVID) for Observational Studies using the Scottish Intercollegiate Guidelines Network Observational Study MEDLINE Search Filter (available at: <http://www.sign.ac.uk/methodology/filters.html>)

1. epidemiologic studies/
2. exp case control studies/
3. exp Cohort Studies/
4. case control.tw.
5. (cohort adj (study or studies)).tw.
6. cohort analy\$.tw.
7. (follow up adj (study or studies)).tw.
8. (observational adj (study or studies)).tw.
9. longitudinal.tw.
10. retrospective.tw.
11. cross sectional.tw.
12. Cross-Sectional Studies/
13. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12
14. (alacepril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril).mp.
15. (moveltipril or pentopril or perindopril or quinapril or ramipril or spirapril or temocapril or teprotide or trandolapril or zofenopril).mp.
16. (losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan).mp.
17. Angiotensin-Converting Enzyme Inhibitors/
18. Angiotensin II Type 1 Receptor Blockers/
19. (ACEI or ARB).mp.
20. 14 or 15 or 16 or 17 or 18 or 19
21. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
- 22. 13 and 20 and 21**

Appendix A: Search Strategies

EMBASE (Silver Platter) for Observational Studies using the Scottish Intercollegiate Guidelines Network Observational Study EMBASE Search Filter (available at: <http://www.sign.ac.uk/methodology/filters.html>)

1. Clinical study/
2. Case control study
3. Family study/
4. Longitudinal study/
5. Retrospective study/
6. Prospective study/
7. Randomized controlled trials/
8. 6 not 7
9. Cohort analysis/
10. (Cohort adj (study or studies)).mp.
11. (Case control adj (study or studies)).tw.
12. (follow up adj (study or studies)).tw.
13. (observational adj (study or studies)).tw.
14. (epidemiologic\$ adj (study or studies)).tw.
15. (cross sectional adj (study or studies)).tw.
16. Or/1-5,8-15
17. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
- 18. 16 AND 17**

Appendix A: Search Strategies

MEDLINE (OVID) for Systematic Reviews Using the Cochrane Highly Sensitive and Specific Search Strategy (Sensitivity and Precision Maximizing Version 2008)

1. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
2. (alacepril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril).mp.
3. (moveltipril or pentopril or perindopril or quinapril or ramipril or spirapril or temocapril or teprotide or trandolapril or zofenopril).mp.
4. (losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan).mp.
5. 2 or 3 or 4
6. (angiotensin-converting enzyme inhibitors or angiotensin II type 2 receptor blockers or ACEI or ARB).mp.
7. 5 or 6
8. (coronary artery disease or coronary disease or myocardial ischemia or angina pectoris or unstable angina or arterial occlusive diseases or peripheral vascular disease or vascular disease or atherosclerosis or cardiovascular diseases or carotid artery diseases).mp.
9. 1 or 8
10. 7 or 9
11. meta-analysis.pt.
12. search.tw.
13. cochrane database of systematic reviews.jn.
14. medline or systematic review.tw.
15. 11 or 12 or 13 or 14
- 16. 10 AND 15**

Appendix A: Search Strategies

Cochrane Database of Systematic Reviews (OVID) for Systematic Reviews

1. (((preserved adj left) or (stable adj cad) or (stable adj chd) or (stable adj coronary) or (preserved adj coronary) or (preserved adj systolic) or (preserved adj ventricular) or (preserved adj lvef) or (preserved adj ef) or (preserved adj ejection)) or (intact adj left) or (intact adj systolic) or (intact adj ventricular) or (intact adj lvef) or (intact adj ef) or (normal adj systolic) or (normal adj ventricular) or (normal adj lvef) or (normal adj ef)).mp
2. (alacepril or benazepril or captopril or ceronapril or cilazapril or delapril or enalapril or fosinopril or imidapril or libenzapril or lisinopril or moexipril).mp.
3. (moveltipril or pentopril or perindopril or quinapril or ramipril or spirapril or temocapril or teprotide ortrandolapril or zofenopril).mp.
4. (losartan or olmesartan or telmisartan or valsartan or eprosartan or candesartan or tasosartan or irbesartan).mp.
5. 2 or 3 or 4
6. (angiotensin-converting enzyme inhibitors or angiotensin II type 2 receptor blockers or ACEI or ARB).mp.
7. 5 or 6
8. (coronary artery disease or coronary disease or myocardial ischemia or angina pectoris or unstable angina or arterial occlusive diseases or peripheral vascular disease or vascular disease or atherosclerosis or cardiovascular diseases or carotid artery diseases).mp.
9. 1 or 8
- 10. 7 AND 9**

Appendix B: List of Excluded Studies

Efficacy/Harms Search

1. Ahmad J, Siddiqui MA, Ahmad H. Effective postponement of diabetic nephropathy with enalapril in normotensive type 2 diabetic patients with microalbuminuria. *Diabetes Care* 1997;20(10):1576-58.
2. Aksnes TA, Kjeldsen SE, Rostup M, et al. Impact of new-onset diabetes mellitus on cardiac outcomes in the Valsartan Antihypertensive Long-term Use Evaluation (VALUE) trial population. *Hypertension* 2007;50(3):467-73.
3. Alderman EL, Botas J. Selection of revascularization for patients with stable angina pectoris. *Coron Artery Dis* 1993;4(12):1061-7.
4. Al-Khadra AS, Salem DN, Rand WM, et al. Antiplatelet agents and survival: a cohort analysis from the Studies of Left Ventricular Dysfunction (SOLVD) trial. *J Am Coll Cardiol* 1998;31(2):419-25.
5. Al-Khadra AS, Salem DN, Rand WM, et al. Antiplatelet agents and survival: a cohort analysis from the Studies of Left Ventricular Dysfunction (SOLVD) trial. *J Am Coll Cardiol* 1998;31(2):419-25.
6. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288(23):2981-97.
7. Ambrosioni E, Borghi C, Magnani B. Early treatment of acute myocardial infarction with angiotensin-converting enzyme inhibition: safety considerations. SMILE pilot study working party. *Am J Cardiol* 1991;68(14):101D-110D.
8. Anavekar NS, McMurray JJ, Velazquez EJ, et al. Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med* 2004;351(13):1285-95.
9. Anavekar NS, Solomon SD, McMurray JD, et al. Comparison of renal function and cardiovascular risk following acute myocardial infarction in patients with and without diabetes mellitus. *Am J Cardiol* 2008;101(7):925-9.
10. Anderson C. Rationale and design of the cardiac magnetic resonance imaging substudy of The ONTARGET Trial Programme. *J Int Med Res* 2005;33(Suppl 1):50A-57A.
11. Anderson TJ, Elstein E, Haber H, et al. Comparative study of ACE-inhibition, angiotensin II antagonism, and calcium channel blockade on flow-mediated vasodilation in patients with coronary disease (BANFF study). *J Am Coll Cardiol* 2000;35(1):60-6.
12. Arima H, Hart RG, Colman S, et al. Perindopril-based blood pressure-lowering reduces major vascular events in patients with atrial fibrillation and prior stroke or transient ischemic attack. *Stroke* 2005;36(10):2164-9.
13. Arima H, Tzourio C, Butcher K, et al. Prior events predict cerebrovascular and coronary outcomes in the PROGRESS trial. *Stroke* 2006;37(6):1497-1502.
14. Arnett DK, Davis BR, Ford CE, et al. Pharmacogenetic association of the angiotensin-converting enzyme insertion/deletion polymorphism on blood pressure and cardiovascular risk in relation to antihypertensive treatment: the Genetics of Hypertension-Associated Treatment (GenHAT) study. *Circulation* 2005;111(25):3374-83.

Appendix B: List of Excluded Studies

15. Asselbergs FW, Diercks GF, Hillege HL, et al. Effects of fosinopril and pravastatin on cardiovascular events in subjects with microalbuminuria. *Circulation* 2004;110(18):2809-16.
16. Asselbergs FW, Hillege HL, van Gilst WH. Framingham score and microalbuminuria: combined future targets for primary prevention? *Kidney Int* 2004;6(Suppl 92):S111-4.
17. Athyros VG, Mikhailidis DP, Papageorgiou AA, et al. Effect of statins and ACE inhibitors alone and in combination on clinical outcome in patients with coronary heart disease. *J Hum Hypertens* 2004;18(11):781-8.
18. Atthobari J, Asselbergs FW, Boersma C, et al. Cost-effectiveness of screening for albuminuria with subsequent fosinopril treatment to prevent cardiovascular events: A pharmacoeconomic analysis linked to the prevention of renal and vascular endstage disease (PREVEND) study and the prevention of renal and vascular endstage disease intervention trial (PREVEND IT). *Clin Ther* 2006;28(3):432-44.
19. Atthobari J, Brantsma AH, Gansevoort RT, et al. The effect of statins on urinary albumin excretion and glomerular filtration rate: results from both a randomized clinical trial and an observational cohort study. *Nephrol Dial Transplant* 2006;21:3106-14.
20. Baba S, and the J-MIND Study Group. Nifedipine and enalapril equally reduce the progression of nephropathy in hypertensive type 2 diabetics. *Diabetes Res Clin Pract* 2001;54(3):191-201.
21. Bakins GL. Benefits of combination therapy for achieving goal blood pressure in high CV risk patients. *Cardiovasc J S Afr* 2001;12:54-5.
22. Bakris GL, Ruilope L, Locatelli F, et al. Treatment of microalbuminuria in hypertensive subjects with elevated cardiovascular risk: results of the IMPROVE trial. *Kidney Int* 2007;72:879-85.
23. Barzilay JI, Jones CL, Davis BR. Baseline characteristics of the diabetic participants in the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Diabetes Care* 2001;24(4):654-8.
24. Baumgart P. [Antihypertensive therapy: risk stratification in diabetes and cardiac diseases.] *MMW Fortschr Med* 2006;148(11):57-60. (German).
25. Bayliss J, Canepa-Anson R, Norell M, et al. The renal response to neuroendocrine inhibition in chronic heart failure: double-blind comparison of captopril and prazosin. *Eur Heart J* 1986;7(10):877-84.
26. Berger PB, Holmes DR, Ohman EM, et al. Restenosis, reocclusion and adverse cardiovascular events after successful balloon angioplasty of occluded versus nonoccluded coronary arteries. Results from the Multicenter American Research Trial With Cilazapril After Angioplasty to Prevent Transluminal Coronary Obstruction and Restenosis (MARCATOR). *J Am Coll Cardiol* 1996;27(1):1-7.
27. Berl T, Hunsicker LG, Lewis JB, et al. Cardiovascular outcomes in the irbesartan diabetic nephropathy trial of patients with type 2 diabetes and overt nephropathy. *Ann Intern Med* 2003;138(7):542-9.
28. Berl T, Hunsicker LG, Lewis JB, et al. Impact of achieved blood pressure on cardiovascular outcomes in the Irbesartan Diabetic Nephropathy Trial. *J Am Soc Nephrol* 2005;16(7):2170-9.
29. Biasucci LM, Lombardi M, Piro M, et al. Irbesartan significantly reduces C reactive protein concentrations after 1 month of treatment in unstable angina. *Heart* 2005;91(5):670-1

Appendix B: List of Excluded Studies

30. Bibbins-Domingo K, Lin F, Vittinghoff E, et al. Renal insufficiency as an independent predictor of mortality among women with heart failure. *J Am Coll Cardiol* 2004;44(8):1593-1600.
31. Bjorholt I, Andersson FL, Kahan T, et al. The cost-effectiveness of ramipril in the treatment of patients at high risk of cardiovascular events: a Swedish sub-study to the HOPE study. *J Intern Med* 2002;251(6):508-17.
32. Black HR, Davis B, Barzilay J, et al. Metabolic and clinical outcomes in nondiabetic individuals with the metabolic syndrome assigned to chlorthalidone, amlodipine, or lisinopril as initial treatment for hypertension: a report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Diabetes Care* 2008;31(2):353-60.
33. Blankenberg S, McQueen MJ, Smieja M, et al. Comparative impact of multiple biomarkers and N-Terminal pro-brain natriuretic peptide in the context of conventional risk factors for the prediction of recurrent cardiovascular events in the Heart Outcomes Prevention Evaluation (HOPE) Study. *Circulation* 2005;114(3):201-8.
34. Boersma C, Carides GW, Atthobari J, et al. An economic assessment of losartan-based versus atenolol-based therapy in patients with hypertension and left-ventricular hypertrophy: results from the Losartan Intervention For Endpoint reduction (LIFE) study adapted to The Netherlands. *Clin Ther* 2007;29(5):963-71.
35. Bohm M, Baumhakel M, Probstfield JL, et al. Sexual function, satisfaction, and association of erectile dysfunction with cardiovascular disease and risk factors in cardiovascular high-risk patients: substudy of the ONgoing Telmisartan Alone and in Combination with Ramipril Global Endpoint Trial/Telmisartan Randomized AssessmentNT Study in ACE-INtolerant Subjects with Cardiovascular Disease (ONTARGET/TRANSCEND). *Am Heart J* 2007;154(1):94-101.
36. Boldt J, Rothe G, Schindler E, et al. Can clonidine, enoximone, and enalaprilat help to protect the myocardium against ischaemia in cardiac surgery? *Heart* 1996;76(3):207-13.
37. Boner G, Cooper ME, McCarroll K, et al. Adverse effects of left ventricular hypertrophy in the reduction of endpoints in NIDDM with the angiotensin II antagonist losartan (RENAAL) study. *Diabetologia* 2005;48(10):1980-7.
38. Bots ML, Remme WJ, Luscher TF, et al. ACE inhibition and endothelial function: main findings of PERFECT, a sub-study of the EUROPA trial. *Cardiovasc Drugs Ther* 2007;21(4):269-79.
39. Bots ML, Remme WJ, Luscher TF, et al. PERindopril-Function of the Endothelium in Coronary Artery Disease Trial: The PERFECT Study-Sub Study of EUROPA: Rationale and Design. *Cardiovasc Drugs Ther* 2002;16(3):227-36.
40. Boulanger JM, Hill MD. Morbidity and mortality after stroke--eprosartan compared with nitrendipine for secondary prevention: principal results of a prospective randomized controlled study (MOSES). *Stroke* 2006;37(2):335-6.
41. Brener SJ, Ivanc TB, Poliszczuk R, et al. Antihypertensive therapy and regression of coronary artery disease: insights from the Comparison of Amlodipine versus Enalapril to Limit Occurrences of Thrombosis (CAMELOT) and Norvasc for Regression of Manifest Atherosclerotic Lesions by Intravascular Sonographic Evaluation (NORMALISE) trials. *Am Heart J* 2006;152(6):1059-63.

Appendix B: List of Excluded Studies

42. Brenner BM, Cooper ME, Zeeuw DD, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345(12):861-9.
43. Briggs A, Mihaylova B, Sculpher M, et al. Cost effectiveness of perindopril in reducing cardiovascular events in patients with stable coronary artery disease using data from the EUROPA study. *Heart* 2007;93(9):1081-6.
44. Burduli FY, Khadzhidis PK, Vatsadze TG, et al. [Use of prazosin and capoten in the treatment of heart failure in patients with ischemic heart disease.] *Kardiologia* 1989;29:49-52. (Russian).
45. Campbell DJ, Woodward M, Chalmers JP, et al. Perindopril-based blood pressure-lowering therapy reduces amino-terminal-pro-B-type natriuretic peptide in individuals with cerebrovascular disease. *J Hypertens* 2007;25(3):699-705.
46. Capri S, Perlini S. Cost-effectiveness in Italy of preventive treatment with ramipril in patients at high risk of cardiovascular events. *Curr Med Res Opin* 2005;21(6):913-21.
47. Carson P, Massie BM, McKelvie R, et al. The Irbesartan in Heart Failure with Preserved Systolic Function. *J Card Fail* 2005;11(8):576-85.
48. Cashin-Hemphill L, Holmvang G, Chan RC, et al. Angiotensin-converting enzyme inhibition as antiatherosclerotic therapy: no answer yet. QUIET Investigators. QUinapril Ischemic Event Trial. *Am J Cardiol* 1999;83(1):43-7.
49. Cashin-Hemphill L, Holmvang G, Chan RC, et al. Angiotensin-converting enzyme inhibition as antiatherosclerotic therapy: no answer yet. QUIET Investigators. QUinapril Ischemic Event Trial. *Am J Cardiol* 1999;83(1):43-7.
50. Catalano M, Libretti A. Captopril for the treatment of patients with hypertension and peripheral vascular disease. *Angiology* 1985;36(5):293-6.
51. Ceconi C, Fox KM, Remme W, et al. ACE inhibition with perindopril and endothelial function. Results of a substudy of the EUROPA study: PERTINENT. *Cardiovasc Res* 2007;73(1):237-46.
52. Cesari M, Kritchevsky SB, Atkinson HH, et al. Angiotensin-converting enzyme inhibition and novel cardiovascular risk biomarkers: results from the Trial of Angiotensin Converting Enzyme Inhibition and Novel cardiovascular risk factors (TRAIN) study. *Am Heart J* 2009;157(2):e1-e8.
53. Chaitman BR, Ivleva AY, Ujda M, et al. Antianginal efficacy of omapatrilat in patients with chronic angina pectoris. *Am J Cardiol* 2005;95(11):1283-9.
54. Charbonneau F. Background rationale for a study to assess normalization of brachial artery forearm flow function. *Can J Cardiol* 1998;14(Suppl D):16D-17D.
55. Coca A, Messerli FH, Benetos A, et al. Predicting stroke risk in hypertensive patients with coronary artery disease: a report from the INVEST trial. *Stroke* 2008;39(2):343-8.
56. Cohen-Solal A, McMurray JJ, Swedberg K, et al. for the CHARM Investigators. Benefits and safety of candesartan treatment in heart failure are independent of age: insights from the Candesartan in Heart Failure – Assessment of Reduction in Mortality and morbidity programme. *Eur Heart J* 2008;29(24):3022-8.
57. Colombo GL, Caruggi M, Ottolini C, et al. Candesartan in Heart failure: Assessment of Reduction in Mortality and morbidity (CHARM) and resource utilization and costs in Italy. *Vasc Health Risk Manag* 2008;4(1):223-34.
58. Cooper DeHoff RM, Handberg EM, Cohen J, et al. Characteristics of contemporary patients with hypertension and coronary artery disease. *Clin Cardiol* 2004;27(10):571-6.

Appendix B: List of Excluded Studies

59. Dagenais GR, Yi Q, Lonn E, et al. Impact of cigarette smoking in high-risk patients participating in a clinical trial. A substudy from the Heart Outcomes Prevention Evaluation (HOPE) trial. *Eur J Cardiovasc Prev Rehabil* 2005;12(1):75-81.
60. Dahlof B, Devereux RB, Kjeldsen SE, et al. Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002;359:995-1003.
61. Daly CA, Hildebrandt P, Bertrand M, et al. Adverse prognosis associated with the metabolic syndrome in established coronary artery disease: data from the EUROPA trial. *Heart* 2007;93(11):1406-11.
62. Dauterman KW, Go AS, Rowell R, et al. Congestive heart failure with preserved systolic function in a statewide sample of community hospitals. *J Card Fail* 2001;7(3):221-8.
63. Dauterman KW, Go AS, Rowell R, et al. Congestive heart failure with preserved systolic function in a statewide sample of community hospitals. *J Card Fail* 2001;7(3):221-8.
64. Davis BR, Arnett DK, Boerwinkle E, et al. Antihypertensive therapy, the alpha-adducin polymorphism, and cardiovascular disease in high-risk hypertensive persons: the Genetics of Hypertension-Associated Treatment Study. *The Pharmacogenomics Journal* 2007;7(2):112-22.
65. Davis BR, Cutler JA, Gordon D, et al. Rationale and design for the Antihypertensive and Lipid Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). ALLHAT Research Group. *Am J Hypertens* 1996;9(4 pt 1):342-60.
66. Davis BR, Kostis JB, Simpson LM, et al. for the ALLHAT Collaborative Research Group. Heart failure with preserved and reduced left ventricular ejection fraction in the antihypertensive and lipid-lowering treatment to prevent heart attack trial. *Circulation* 2008;118(22):2259-67.
67. de Simone G, Wachtell K, Palmieri V, et al. Body build and risk of cardiovascular events in hypertension and left ventricular hypertrophy: the LIFE (Losartan Intervention For Endpoint reduction in hypertension) study. *Circulation* 2005;111(15):1924-31.
68. de Zeeuw D, Remuzzi G, Parving HH, et al. Albuminuria, a therapeutic target for cardiovascular protection in type 2 diabetic patients with nephropathy. *Circulation* 2004;110(8):921-7.
69. Desmet W, Vrolix M, Scheerder ID, et al. Angiotensin-converting enzyme inhibition with fosinopril sodium in the prevention of restenosis after coronary angioplasty. *Circulation* 1994;89:385-92.
70. Devereux RB, Dahlof B, Kjeldsen SE, et al. Effects of losartan or atenolol in hypertensive patients without clinically evident vascular disease: a substudy of the LIFE randomized trial. *Ann Intern Med* 2003;139(3):169-77.
71. Devereux RB, Wachtell K, Gerds E, et al. Prognostic significance of left ventricular mass change during treatment of hypertension. *JAMA* 2004;292(19):2350-6.
72. Diercks GF, Janssen WM, van Boven AJ, et al. Rationale, design, and baseline characteristics of a trial of prevention of cardiovascular and renal disease with fosinopril and pravastatin in nonhypertensive, nonhypercholesterolemic subjects with microalbuminuria (the Prevention of RENal and Vascular ENdstage Disease Intervention Trial [PREVEND IT]). *Am J Cardiol* 2000;86(6):635-8.
73. Domanski MJ, Jablonski KA, Rice MM, et al. Obesity and cardiovascular events in patients with established coronary disease. *Eur Heart J* 2006;27(12):1416-22.

Appendix B: List of Excluded Studies

74. Duprez DA, Florea ND, Jones K, et al. Beneficial effects of valsartan in asymptomatic individuals with vascular or cardiac abnormalities: the DETECTIV Pilot Study. *J Am Coll Cardiol* 2007;50(9):835-9.
75. Ekbom T, Linjer E, Hedner T, et al. Cardiovascular events in elderly patients with isolated systolic hypertension. A subgroup analysis of treatment strategies in STOP-Hypertension-2. *Blood Press* 2004;13(3):137-41.
76. Ekeberg O, Klemsdal TO, Kjeldsen SE. Quality of life on enalapril after acute myocardial infarction. *Eur Heart J* 1994;15(8):1135-9.
77. Erinc K, Yamani MH, Starling RC, et al. The effect of combined Angiotensin-converting enzyme inhibition and calcium antagonism on allograft coronary vasculopathy validated by intravascular ultrasound. *J Heart Lung Transplant* 2005;24(8):1033-8.
78. Estacio RO, Coll JR, Tran ZV, et al. Effect of intensive blood pressure control with valsartan on urinary albumin excretion in normotensive patients with type 2 diabetes. *Am J Hypertens* 2006;19(12):1241-8.
79. Estacio RO, Jeffers BW, Gifford N, et al. Effect of blood pressure control on diabetic microvascular complications in patients with hypertension and type 2 diabetes. *Diabetes Care* 2000;23(Suppl 2):B54-64.
80. Estacio RO, Jeffers BW, Hiatt WR, et al. The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med* 1998;338:645-52.
81. Evidence-based Clinical Practice. Addition of an ACE inhibitor to conventional treatment is not beneficial in people with stable coronary artery disease and preserved left ventricular function. *Evid Based Healthc Public Health* 2005;9(3):259-60.
82. Ezekowitz JA, Lee DS, Tu JV, et al. Comparison of one-year outcome (death and rehospitalization) in hospitalized heart failure patients with left ventricular ejection fraction >50% versus those with ejection fraction <50%. *Am J Cardiol* 2008;102(1):79-83.
83. Fagard RH, Staessen JA, Thijs L, et al. Prognostic significance of electrocardiographic voltages and their serial changes in elderly with systolic hypertension. *Hypertension* 2004;44(4):459-64.
84. Fallowfield JM, Blenkinsopp J, Raza A, et al. Post-marketing surveillance of lisinopril in general practice in the UK. *Br J Clin Pract* 1993;47(6):296-304.
85. Ferrari R, and The PREAMI Investigators. Effects of angiotensin-converting enzyme inhibition with perindopril on left ventricular remodeling and clinical outcome: results of the randomized Perindopril and Remodeling in Elderly with Acute Myocardial Infarction (PREAMI) Study. *Arch Intern Med* 2006;166(6):659-66.
86. Fossum E, Gleim GW, Kjeldsen SE, et al. The effect of baseline physical activity on cardiovascular outcomes and new-onset diabetes in patients treated for hypertension and left ventricular hypertrophy: the LIFE study. *J Intern Med* 2007;262(4):439-48.
87. Fukami K, Hiromori K, Kishida H, et al. [Assessment and controversy of long-term medical treatment in ischemic heart disease. Japanese Angina and Myocardial infarction Prospective (JAMP) study.] *Jpn Circ J* 1994;58 Suppl 4:1293-6. (Japanese).
88. Gaede PH, Jepsen PV, Larsen JN, et al. [The Steno-2 study. Intensive multifactorial intervention reduces the occurrence of cardiovascular disease in patients with type 2 diabetes.] *Ugeskr Laeger* 2003;165(26):2658-61. (Danish).

Appendix B: List of Excluded Studies

89. Geluk CA, Asselbergs FW, Hillege HL, et al. Impact of statins in microalbuminuric subjects with the metabolic syndrome: a substudy of the PREVEND Intervention Trial. *Eur Heart J* 2005;26(13):1314-20.
90. Gerds E, Wachtell K, Omvik P, et al. Left atrial size and risk of major cardiovascular events during antihypertensive treatment: losartan intervention for endpoint reduction in hypertension trial. *Hypertension* 2007;49(2):311-6.
91. Gerstein HC, Pogue J, Mann JF, et al. The relationship between dysglycaemia and cardiovascular and renal risk in diabetic and non-diabetic participants in the HOPE study: a prospective epidemiological analysis. *Diabetologia* 2005;48(9):1749-55.
92. Gerstein HC, Swedberg K, Carlsson J, et al. for the CHARM Program Investigators. The hemoglobin A1c level as a progressive risk factor for cardiovascular death, hospitalization for heart failure, or death in patients with chronic heart failure: an analysis of the Candesartan in Heart failure: Assessment of Reduction in Mortality and Morbidity (CHARM) program. *Arch Intern Med* 2008;168(15):1699-704.
93. Gibas M, Mischczak-Smialek J, Madry E. Influence of preventive therapy with quinapril on IL-6 level in patients with chronic stable angina. *Pharmacol Rep* 2007;59(3):330-8.
94. Gnechchi M, Ardissino D. [Effects of ACE inhibitors on angiographic restenosis after coronary stenting (PARIS): double-blind randomized trial.] *Ital Heart J* 2001;2(9):1027-8. (Italian).
95. Go AS, Iribarren C, Chandra M, et al. Statin and beta-blocker therapy and the initial presentation of coronary heart disease. *Ann Intern Med* 2006;144(4):229-38.
96. Hager WD, Davis BR, Riba A, et al. Absence of a deleterious effect of calcium channel blockers in patients with left ventricular dysfunction after myocardial infarction: The SAVE Study Experience. SAVE Investigators. Survival and Ventricular Enlargement. *Am Heart J* 1998;135(3):406-13.
97. Hansson L. The Captopril Prevention Project (CAPPP): description and status. *Am J Hypertens* 1994;7(9 Pt 2):82S-83S.
98. Hansson L, Lithell H, Skoog I, et al. Study on COgnition and Prognosis in the Elderly (SCOPE). *Blood Press* 1999;8(3):177-83.
99. Hansson L, Lithell H, Skoog I, et al. Study on COgnition and Prognosis in the Elderly (SCOPE): baseline characteristics. *Blood Press* 2000;9:146-51.
100. Harrap SB, Tzourio C, Cambien F, et al. The ACE gene I/D polymorphism is not associated with the blood pressure and cardiovascular benefits of ACE inhibition. *Hypertension* 2003;42(3):297-303.
101. Hart WM, Rubio-Terres C, Margalet Fernandez I, et al. [Cost-effectiveness analysis of Ramipril treatment of patients at high-risk of cardiovascular events in Spain.] *An Med Interna* 2002;19(10):515-20 (Spanish).
102. Hawkins NM, Wang D, McMurray JJ, et al. Prevalence and prognostic impact of bundle branch block in patients with heart failure: evidence from the CHARM programme. *Eur J Heart Fail* 2007;9(5):510-7.
103. Hermans WR, Foley DP, Rensing BJ, et al. Morphologic changes during follow-up after successful percutaneous transluminal coronary balloon angioplasty: quantitative angiographic analysis in 778 lesions--further evidence for the restenosis paradox. MERCATOR Study Group (Multicenter European Research trial with Cilazapril after Angioplasty to prevent Transluminal Coronary Obstruction and Restenosis). *Am Heart J* 1994;127(3):483-94.

Appendix B: List of Excluded Studies

104. Hermans WR, Rensing BJ, Foley DP, et al. Therapeutic dissection after successful coronary balloon angioplasty: no influence on restenosis or on clinical outcome in 693 patients. The MERCATOR Study Group (Multicenter European Research Trial with Cilazapril after Angioplasty to prevent Transluminal Coronary Obstruction and Restenosis). *J Am Coll Cardiol* 1992;20(4):767-80.
105. Hermans WR, Rensing BJ, Foley DP, et al. Patient, lesion, and procedural variables as risk factors for luminal re-narrowing after successful coronary angioplasty: a quantitative analysis in 653 patients with 778 lesions. Multicenter European Research Trial with Cilazapril after Angioplasty to prevent Transluminal Coronary Obstruction and Restenosis (MERCATOR) Study Group. *J Cardiovasc Pharmacol* 1993;22(Suppl 4):S45-57.
106. Heyndrickx GR, and the MERCATOR Study Group. Angiotensin-converting enzyme inhibitor in a human model of restenosis. MERCATOR ("Multicenter European Research Trial with Cilazapril after Angioplasty to Prevent Transluminal Coronary Obstruction and Restenosis") Study Group. *Basic Res Cardiol* 1993;88(Suppl 1):169-82.
107. Hofmann M, Bauer R, Handrock R, et al. Prognostic value of the QRS duration in patients with heart failure: a subgroup analysis from 24 centers of Val-HeFT. *J Card Fail* 2005;11(7):523-8.
108. Hoiegggen A, Alderman MH, Kjeldsen SE, et al. The impact of serum uric acid on cardiovascular outcomes in the LIFE study. *Kidney Int* 2004;65(3):1041-9.
109. Hoogwerf BJ, Young JB. The HOPE study. Ramipril lowered cardiovascular risk, but vitamin E did not. *Cleve Clin J Med* 2000;67(4):287-93.
110. Hosomi N, Mizushige K, Ohyama H, et al. Angiotensin-converting enzyme inhibition with enalapril slows progressive intima-media thickening of the common carotid artery in patients with non-insulin-dependent diabetes mellitus. *Stroke* 2001;32(7):1539-45.
111. Hsia J, Jablonski KA, Rice MM, et al. Sudden Cardiac Death in Patients With Stable Coronary Artery Disease and Preserved Left Ventricular Systolic Function. *Am J Cardiol* 2008;101(4):457-61.
112. Hughes S. EUROPA study results: Perindopril benefits broad range of patients with stable coronary disease. *Br J Cardiol* 2003;10(5):339-40.
113. Ibsen, H.; Olsen, M. H.; Wachtell, K. Reduction in albuminuria translates to reduction in cardiovascular events in hypertensive patients: losartan intervention for endpoint reduction in hypertension study. *Hypertension* 2005;45(2):198-202.
114. Ibsen H, Olsen MH, Wachtell K, et al. Does albuminuria predict cardiovascular outcomes on treatment with losartan versus atenolol in patients with diabetes, hypertension, and left ventricular hypertrophy? The LIFE study. *Diabetes Care* 2006;29(3):595-600.
115. Ibsen H, Pedersen OL, Dahlof B, et al. [The effect of losartan versus atenolol on cardiovascular morbidity and mortality in patients with hypertension and ECG-verified left ventricular hypertrophy in the LIFE-study.] *Ugeskr Laeger* 2003;165(5):456-9. (Danish).
116. Ikeda N, Nishimura S, Kyo S, et al. Valsartan cardio-renal protection in patients undergoing coronary angiography complicated with chronic renal insufficiency (VAL-CARP) trial: rationale and design. *Circ J* 2006;70(5):548-52.
117. Ishani A, Herzog CA, Collins AJ, et al. Cardiac medications and their association with cardiovascular events in incident dialysis patients: cause or effect?. *Kidney Int* 2004;65(3):1017-25.

Appendix B: List of Excluded Studies

118. Janardhanan R, Kenchaiah S, Velazquez EJ, et al. Extent of coronary artery disease as a predictor of outcomes in acute myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *Am Heart J* 2006;152(1):183-9.
119. Jezek T, Balazovjech I. The Valsartan Antihypertensive Long-term Use Evaluation (VALUE) trial in Slovakia. *Bratisl Lek Listy* 2003;104(1):19-25
120. Johnstone D, Limacher M, Rousseau M, et al. Clinical characteristics of patients in studies of left ventricular dysfunction (SOLVD). *Am J cardiol* 1992;70(9):894-900.
121. Jonsson B, Buxton M, Hertzman P, et al. Health economics of prevention of coronary heart disease and vascular events: a cost-effectiveness analysis based on the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT). *J Hum Hypertens* 2001;15(Suppl 1):S53-6.
122. Jose P, Skali H, Anavekar N, et al. Increase in creatinine and cardiovascular risk in patients with systolic dysfunction after myocardial infarction. *J Am Soc Nephrol* 2006;17(10):2886-91.
123. Julius S, Kjeldsen SE, Brunner H, et al. VALUE trial: Long-term blood pressure trends in 13,449 patients with hypertension and high cardiovascular risk. *Am J Hypertens* 2003;16(7):544-8.
124. Julius S, Kjeldsen SE, Weber M, et al. Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomised trial. *Lancet* 2004;363(9426):2022-31.
125. Just H, Drexler H, Taylor SH, et al. Captopril versus digoxin in patients with coronary artery disease and mild heart failure. A prospective, double-blind, placebo-controlled multicenter study. The CADS Study Group. *Herz* 1993;18(Suppl 1):S436-43.
126. Kaul U, Chandra S, Bahl VK, et al. Enalapril for prevention of restenosis after coronary angioplasty. *Indian Heart J* 1993;45(6):469-73.
127. Kenchaiah S, Davis BR, Braunwald E, et al. Antecedent hypertension and the effect of captopril on the risk of adverse cardiovascular outcomes after acute myocardial infarction with left ventricular systolic dysfunction: Insights from the Survival and Ventricular Enlargement Trial. *Am Heart J* 2004;148(2):356-64.
128. Kenchaiah S, Pocock SJ, Wang D, et al. Body mass index and prognosis in patients with chronic heart failure: insights from the Candesartan in Heart failure: Assessment of Reduction in Mortality and morbidity (CHARM) program. *Circulation* 2007;116(6):627-36
129. Kessler M, Zannad F, Lehert P, et al. Predictors of cardiovascular events in patients with end-stage renal disease: an analysis from the Fosinopril in dialysis study. *Nephrol Dial Transplant* 2007;22(12):3573-9.
130. Kjeldsen SE, Dahlof B, Devereux RB, et al. Effects of losartan on cardiovascular morbidity and mortality in patients with isolated systolic hypertension and left ventricular hypertrophy: a Losartan Intervention for Endpoint Reduction (LIFE) substudy. *JAMA* 2002;288(12):1491-8.
131. Kjeldsen SE, Julius S, Brunner H, et al. Characteristics of 15,314 hypertensive patients at high coronary risk. The VALUE trial. The Valsartan Antihypertensive Long-term Use Evaluation. *Blood Press* 2001;10(2):83-91.
132. Kleber FX, Niemoller L, Fischer M, et al. Influence of severity of heart failure on the efficacy of angiotensin-converting enzyme inhibition. *Am J Cardiol* 1991;68(14):121D-126D.

Appendix B: List of Excluded Studies

133. Kober L, Torp-Pedersen C, Carlsen JE, et al. A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. Trandolapril Cardiac Evaluation (TRACE) Study Group. *N Engl J Med* 1995;333(25):1670-6.
134. Kontopoulos AG, Athyros VG, Papageorgiou AA, et al. Effect of angiotensin-converting enzyme inhibitors on the power spectrum of heart rate variability in post-myocardial infarction patients. *Coron Artery Dis* 1997;8:517-24.
135. Kowey PR, Dickson TZ, Zhang Z, et al. Losartan and end-organ protection--lessons from the RENAAL study. *Clin Cardiol* 2005;28(3):136-42.
136. Kristianson K, Fyhrquist F, Devereux RB, et al. An analysis of cholesterol control and statin use in the Losartan Intervention for Endpoint Reduction in Hypertension Study. *Clin Ther* 2003;25(4):1186-99.
137. Krysiak R, Okopien B. Pleiotropic effects of angiotensin-converting enzyme inhibitors in normotensive patients with coronary artery disease. *Pharmacol Rep* 2008;60(4):514-23.
138. Lahoz C, Mostaza JM, Mantilla MT, et al. Achievement of therapeutic goals and utilization of evidence-based cardiovascular therapies in coronary heart disease patients with chronic kidney disease. *Am J Cardiol* 2008;101(8):1098-1102.
139. Lamas GA, Mitchell GF, Flaker GC, et al. Clinical significance of mitral regurgitation after acute myocardial infarction. Survival and Ventricular Enlargement Investigators. *Circulation* 1997;96(3):827-33.
140. Landmesser U, Spiekermann S, Preuss C, et al. Angiotensin II induces endothelial xanthine oxidase activation: role for endothelial dysfunction in patients with coronary disease. *Arterioscler Thromb Vasc Biol* 2007;27(4):943-8.
141. Leenen FH, Nwachuku CE, Black HR, et al. Clinical events in high-risk hypertensive patients randomly assigned to calcium channel blocker versus angiotensin-converting enzyme inhibitor in the antihypertensive and lipid-lowering treatment to prevent heart attack trial. *Hypertension* 2006;48(3):374-84.
142. Leor J, Reicher-Reiss H, Goldbourt U, et al. Aspirin and mortality in patients treated with angiotensin-converting enzyme inhibitors: a cohort study of 11,575 patients with coronary artery disease. *J Am Col Cardiol* 1999;33(7):1920-5.
143. Lievre M, Marre M, Chatellier G, et al. The non-insulin-dependent diabetes, hypertension, microalbuminuria or proteinuria, cardiovascular events, and ramipril (DIABHYCAR) study: design, organization, and patient recruitment. DIABHYCAR Study Group. *Control Clin Trials* 2000;21(4):383-96.
144. Lindholm LH, Ibsen H, Dahlöf B, et al. Cardiovascular morbidity and mortality in patients with diabetes in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002;359(9311):1004-10.
145. Linjer E, Jörnmark J, Hedner T, et al. Predictors for high costs of hospital care in elderly hypertensive patients. *Blood Press* 2006;15(4):245-50.
146. Lithell H, Hansson L, Skoog I, et al. The Study on COgnition and Prognosis in the Elderly (SCOPE); outcomes in patients not receiving add-on therapy after randomization. *J Hypertens* 2004;22(8):1605-12.
147. Lithell H, Hansson L, Skoog I, et al. The Study on Cognition and Prognosis in the Elderly (SCOPE): principal results of a randomized double-blind intervention trial. *J Hypertens* 2003;21(5):875-86.

Appendix B: List of Excluded Studies

148. Lobos Bejarano JM, Del Castillo JC. The angiotensin-converting enzyme inhibitors reduce the morbidity-mortality in patients with a stable coronary artery disease, in the absence of left ventricular dysfunction or heart failure. *FMC* 2007;14(3):184.
149. Lonn E, Shaikhholeslami R, Yi Q, et al. Effects of ramipril on left ventricular mass and function in cardiovascular patients with controlled blood pressure and with preserved left ventricular ejection fraction: A substudy of the Heart Outcomes Prevention Evaluation (HOPE) trial. *J Am Coll Cardiol* 2004;43(12):2200-6.
150. Lonn E, Yusuf S, Dzavik V, et al. Effects of ramipril and vitamin E on atherosclerosis: the study to evaluate carotid ultrasound changes in patients treated with ramipril and vitamin E (SECURE). *Circulation* 2001;103(7):919-25.
151. Lonn EM, Yusuf S, Doris CI, et al. Study design and baseline characteristics of the study to evaluate carotid ultrasound changes in patients treated with ramipril and vitamin E: SECURE. *Am J Cardiol* 1996;78(8):914-9.
152. Luders S, Schrader J, Berger J, et al. The PHARAO study: prevention of hypertension with the angiotensin-converting enzyme inhibitor ramipril in patients with high-normal blood pressure: a prospective, randomized, controlled prevention trial of the German Hypertension League. *J Hypertens* 2008;26(7):1487-96.
153. Luno J, Garcia de Vinuesa S. [Control of arterial pressure in diabetic nephropathy.] *Nefrologia* 2001;21(3):240-5. (Spanish).
154. Lynch, A. I.; Boerwinkle, E.; Davis, B. R. Pharmacogenetic association of the NPPA T2238C genetic variant with cardiovascular disease outcomes in patients with hypertension. *JAMA* 2008;299(3):296-307.
155. Magrini G, Nicolosi GL, Chiariello M, et al. [Rationale, characteristics and study design of PREAMI (Perindopril and Remodelling in the Elderly with Acute Myocardial Infarction) study.] *Ital Heart J* 2005;6 (Suppl 7):14S-23S. (Italian).
156. Majahalme, S. K.; Baruch, L.; Aknay, N. Comparison of treatment benefit and outcome in women versus men with chronic heart failure (from the Valsartan Heart Failure Trial). *Am J Cardiol* 2005;95(4):529-32.
157. Mancini GB. Quinapril ischemic event trial. *Am J Cardiol* 2001;88(11):1348.
158. Management Committee on behalf of the High Blood Pressure Research Council of Australia. Australian comparative outcome trial of angiotensin-converting enzyme inhibitor and diuretic based treatment of hypertension in the elderly (ANBP2): Objectives and protocol. *Clin Exp Pharmacol Physiol* 1997;24(2):188-92.
159. Mancini GBJ, Henry GC, Macaya C, et al. Angiotensin-converting enzyme inhibition with quinapril improves endothelial vasomotor dysfunction in patients with coronary artery disease. *Circulation* 1996;94:258-65.
160. Mann J, Julius S. The Valsartan Antihypertensive Long-term Use Evaluation (VALUE) trial of cardiovascular events in hypertension. Rationale and design. *Blood Press* 1998;7(3):176-83.
161. Mann JF, Gerstein HC, Yi Q, et al. Development of renal disease in people at high cardiovascular risk: results of the HOPE randomized study. *J Am Soc Nephrol* 2003;14(3):641-7.
162. Mann JF, Schmieder RE, McQueen M, et al. for the ONTARGET Investigators. Renal outcomes with telmisartan, ramipril, or both, in people at high vascular risk (the ONTARGET study): a multicentre, randomized, double-blind, controlled trial. *Lancet* 2008;372(9638):547-53.

Appendix B: List of Excluded Studies

163. Marre M, Lievre M, Vasmant D, et al. Determinants of elevated urinary albumin in the 4,937 type 2 diabetic subjects recruited for the DIABHYCAR Study in Western Europe and North Africa. *Diabetes Care* 2000;23(Suppl 2):B40-8.
164. Mason B, Matsuyama J, Jue S. Adherence consistency across treatment regimens. *Diabetes Care* 1994;17(4):347-8.
165. Massie BM, Carson PE, McMurray JJ, et al. Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med* 2008;359(23):2456-67.
166. McCall AL. Reducing CVD risk in type 2 DM. *Current Diabetes Report* 2003;3(5):363-4.
167. McConaghy JR. Is treatment of hypertension with acetylcholine esterase inhibitors (ACEIs) superior to other antihypertensives in preventing significant cardiovascular events and death in patients with type 2 diabetes? *J Fam Pract* 2000;49(11):980.
168. McMurray J, Ostergren J, Pfeffer M, et al. Clinical features and contemporary management of patients with low and preserved ejection fraction heart failure: Baseline characteristics of patients in the Candesartan in Heart failure - Assessment of Reduction in Mortality and morbidity. *Eur J Heart Fail* 2003;5(3):261-70.
169. McMurray J, Ostergren J, Pfeffer M, et al. Clinical features and contemporary management of patients with low and preserved ejection fraction heart failure: baseline characteristics of patients in the Candesartan in Heart failure-Assessment of Reduction in Mortality and morbidity (CHARM) programme. *Eur J Heart Fail* 2003;5(3):261-70.
170. McMurray J, Solomon S, Pieper K, et al. The effect of valsartan, captopril, or both on atherosclerotic events after acute myocardial infarction: an analysis of the Valsartan in Acute Myocardial Infarction Trial (VALIANT). *J Am Coll Cardiol* 2006;47(4):726-33.
171. McMurray JJ, Ostergren J, Swedberg K, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet* 2003;362(9386):767-71.
172. McMurray JJV, Carson PE, Komajda M, et al. Heart failure with preserved ejection fraction: Clinical characteristics of 4133 patients enrolled in the I-PRESERVE trial. *Eur J Heart Fail* 2008;10(2):149-56.
173. Mehler PS, Coll JR, Estacio R, et al. Intensive blood pressure control reduces the risk of cardiovascular events in patients with peripheral arterial disease and type 2 diabetes. *Circulation* 2003;107(5):753-6.
174. Menard J, Patchett AA. Angiotensin-converting enzyme inhibitors. *Adv Protein Chem* 2001;56:13-75.
175. Meredith IT, Alison JF, Zhang FM, et al. Captopril potentiates the effects of nitroglycerin in the coronary vascular bed. *J Am Coll Cardiol* 1993;22(2):581-7.
176. Messerli FH, Grossman E. Doxazosin arm of the ALLHAT study discontinued: how equal are antihypertensive drugs? Antihypertensive and Lipid Lowering Treatment to Prevent Heart Attack Trial. *Curr Hypertens Rep* 2000;2(3):241-2.
177. Messerli FH, Grossman E. CAPPP trial. Captopril Prevention Project. *Lancet* 1999;353(9166):1794.
178. Mitchell GF, Dunlap ME, Warnica W, et al. Long-term trandolapril treatment is associated with reduced aortic stiffness: The prevention of events with angiotensin-converting enzyme inhibition hemodynamic substudy. *Hypertension* 2007;49(6):1271-7.
179. Mitsuyama K. [Multiphasic action of AT1 receptor blockers and their therapeutic effects for cardiovascular diseases.] *Nippon Yakurigaku Zasshi* 2008;131(3):176-7. (Japanese).

Appendix B: List of Excluded Studies

180. Mochizuki S, Dahlof B, Shimizu M, et al. Valsartan in a Japanese population with hypertension and other cardiovascular disease (Jikei Heart Study): a randomised, open-label, blinded endpoint morbidity-mortality study. *Lancet* 2007;369(9571):1431-9.
181. Mochizuki S, Shimizu M, Taniguchi I, et al. JIKEI HEART Study--a morbi-mortality and remodeling study with valsartan in Japanese patients with hypertension and cardiovascular disease. *Cardiovasc Drugs Ther* 2004;18(4):305-9.
182. Nesbitt SD. Perspectives on prehypertension. *J Cardiometab Syndr* 2006;1(5):364-5
183. Nielsen FS. Diabetic nephropathy in non-insulin dependent diabetes mellitus. Cardiovascular risk factors and antihypertensive treatment. *Dan Med Bull* 2000;47(4):249-70.
184. Niskanen L, Hedner T, Hansson L, et al. Reduced cardiovascular morbidity and mortality in hypertensive diabetic patients on first-line therapy with an ACE inhibitor compared with a diuretic/beta-blocker-based treatment regimen: a subanalysis of the Captopril Prevention Project. *Diabetes Care* 2001;24(12):2091-6.
185. Norris K, Bourgoigne J, Gassman J, et al. Cardiovascular outcomes in the African American Study of Kidney Disease and Hypertension (AASK) Trial. *Am J Kidney Dis* 2006;48(5):739-51.
186. Odenthal HJ, Josephs W. [Angiotensin-converting enzyme inhibition and angina pectoris.] *Dtsch Med Wochenschr* 1992;117(48):1849-53. (German).
187. Ogihara T, Nakao K, Fukui T, et al. Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. *Hypertension* 2008;51(2):393-8.
188. Okamura A, Ohishi M, Rakugi H, et al. Pharmacogenetic analysis of the effect of angiotensin-converting enzyme inhibitor on restenosis after percutaneous transluminal coronary angioplasty. *Angiology* 1999;50(10):811-22.
189. Okimoto T, Imazu M, Hayashi Y, et al. Quinapril with high affinity to tissue angiotensin-converting enzyme reduces restenosis after percutaneous transcatheter coronary intervention. *Cardiovasc Drugs Ther* 2001;15(4):323-9.
190. Okumura K, Sone T, Kondo J, et al. Quinapril prevents restenosis after coronary stenting in patients with angiotensin-converting enzyme D allele. *Circ J* 2002;66(4):311-6.
191. Olsen MH, Wachtell K, Dahlof B, et al. The effect of losartan compared with atenolol on the incidence of revascularization in patients with hypertension and electrocardiographic left ventricular hypertrophy. The LIFE study. *J Hum Hypertens* 2006;20(6):460-4.
192. Olsson LG, Swedberg K, Ducharme A, et al. Atrial fibrillation and risk of clinical events in chronic heart failure with and without left ventricular systolic dysfunction: results from the Candesartan in Heart failure-Assessment of Reduction in Mortality and morbidity (CHARM) program. *J Am Coll Cardiol* 2006;47(10):1997-2004.
193. Omland T, Sabatine MS, Jablonski K, et al. Prognostic Value of B-Type Natriuretic Peptides in Patients With Stable Coronary Artery Disease. The PEACE Trial. *J Am Coll Cardiol* 2007;50(3):205-14.
194. Oosterga M, Voors AA, Buikema H, et al. Angiotensin II formation in human vasculature after chronic ACE inhibition: A prospective, randomized, placebo-controlled study. *Cardiovasc Drugs Ther* 2000;14:55-60.
195. Oosterga M, Voors AA, Pinto YM, et al. Effects of quinapril on clinical outcome after coronary artery bypass grafting (The QUO VADIS study). *Am J Cardiol* 2001;87(5):542-6.

Appendix B: List of Excluded Studies

196. Opie L, Dalby T, Naidoo DP. Perindopril (Coversyl) prevents cardiovascular death and MI in coronary disease patients regardless of their cardiovascular risk. *Cardiovasc J S Afr* 2003;14(5):277-9.
197. Os I, Franco V, Kjeldsen SE, et al. Effects of losartan in women with hypertension and left ventricular hypertrophy: results from the Losartan Intervention for Endpoint Reduction in Hypertension Study. *Hypertension* 2008;51(4):1103-8.
198. Ostergren J, Sleight P, Dagenais G, et al. Impact of ramipril in patients with evidence of clinical or subclinical peripheral arterial disease. *Eur Heart J* 2004;25(1):17-24.
199. Ostergren J, Bjorhold I, Andersson F, et al. Pharmacoeconomic impact of HOPE. *Int J Clin Pract* 2001;Suppl 117:19-21.
200. Otsuka M, Yamamoto H, Okimoto T, et al. Long-term effects of quinapril with high affinity for tissue angiotensin-converting enzyme after coronary intervention in Japanese. *Am Heart J* 2004;147(4):662-8.
201. Ottersetted JE, Sleight P. The HOPE study: comparison with other trials of secondary prevention. *Eur Heart J* 2001;22(15):1307-10.
202. Ozdemir M, Arslan U, Turkoglu S, et al. Losartan improves heart rate variability and heart rate turbulence in heart failure due to ischemic cardiomyopathy. *J Card Fail* 2007;13(10):812-7.
203. Palmas W, Ma S, Psaty B, et al. Antihypertensive medications and C-reactive protein in the multi-ethnic study of atherosclerosis. *Am J Hypertens* 2007;20(3):233-41.
204. Passa P, Chatellier G. The DIAB-HYCAR Study. *Diabetologia* 1996;39(12):1662-7.
205. Patel A, Chalmers J, Poulter N. ADVANCE: action in diabetes and vascular disease. *J Hum Hypertens* 2005;19(Suppl 1):S27-S32.
206. Pepine CJ, Rouleau JL, Annis K, et al. Effects of angiotensin-converting enzyme inhibition on transient ischemia: the Quinapril Anti-Ischemia and Symptoms of Angina Reduction (QUASAR) trial. *J Am Coll Cardiol* 2003;42(12):2049-59.
207. Perepech NB. [Ischemic heart disease with low risk of vascular complications -- a novel indication for angiotensin converting enzyme inhibitors (results of EUROPA trial).] *Kardiologija* 2004;44(3):106-10. (Russian).
208. Perkovic V, Ninomiya T, Arima H, et al. Chronic kidney disease, cardiovascular events, and the effects of perindopril-based blood pressure lowering: data from the PROGRESS study. *J Am Soc Nephrol* 2007;18(10):2766-72.
209. Persson H, Lonn E, Edner M, et al. Diastolic dysfunction in heart failure with preserved systolic function: need for objective evidence: results from the CHARM Echocardiographic Substudy-CHARMES. *J Am Coll Cardiol* 2007;49(6):687-94.
210. Peters S, Gotting B, Trummel M, et al. Valsartan for prevention of restenosis after stenting of type B2/C lesions: the VAL-PREST trial. *J Invas Cardiol* 2001;13(2):93-7.
211. Pfeffer MA, Braunwald E, Moye LA, et al. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. Results of the survival and ventricular enlargement trial. The SAVE Investigators. *N Engl J Med* 1992;327(10):669-677.
212. Pfeffer, MA, McMurray JJV, Velazquez EJ, et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;349(20):1893-906.

Appendix B: List of Excluded Studies

213. Pfeffer MA, Swedberg K, Granger CB, et al. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet* 2003;362:759-66.
214. Pfeffer MA, Domanski M, Verter J, et al. The continuation of the prevention of events with angiotensin-converting enzyme inhibition (PEACE) trial. *Am Heart J* 2001;142(3):375-7.
215. Piller LB, Ford CE, Davis BR, et al. Incidence and predictors of angioedema in elderly hypertensive patients at high risk for cardiovascular disease: A report from the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *J Clin Hypertens* 2006 8(9):649-56.
216. Poole-Wilson PA, Lubsen J. Losartan for cardiovascular disease in patients with and without diabetes in the LIFE study. *Lancet* 2002;359:2199.
217. Procopiou M. The ADVANCE trial. *Lancet* 2008;371:25.
218. The PROGRESS Collaborative Group. Effects of a perindopril-based blood pressure lowering regimen on cardiac outcomes among patients with cerebrovascular disease. *Eur Heart J* 2003;24(5):475-84
219. Rahman M, Pressel S, Davis BR, et al. Cardiovascular outcomes in high-risk hypertensive patients stratified by baseline glomerular filtration rate. *Ann Intern Med* 2006;144(3):172-80.
220. Rahman M, Pressel S, Davis BR, et al. Renal outcomes in high-risk hypertensive patients treated with an angiotensin-converting enzyme inhibitor or a calcium channel blocker vs a diuretic: A report from the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *Arch Intern Med* 2005;165:936-46.
221. Rahman M, Pressel S, Davis BR, et al. Cardiovascular outcomes in high-risk hypertensive patients stratified by baseline glomerular filtration rate. *Ann Intern Med* 2006;144(3):172-80.
222. Rayner B. Selective imidazoline agonist moxonidine plus the ACE inhibitor ramipril in hypertensive patients with impaired insulin sensitivity: partners in a successful MARRIAGE? *Curr Med Res Opin* 2004;20(3):359-67.
223. Reims HM, Oparil S, Kjeldsen SE, et al. Losartan benefits over atenolol in non-smoking hypertensive patients with left ventricular hypertrophy: The LIFE study. *Blood Press* 2004;13(6):376-84.
224. Redekop WK, Orlewska E, Maciejewski P, et al. Costs and effects of secondary prevention with perindopril in stable coronary heart disease in Poland: an analysis fo the EUROPA study including 1251 Polish patients. *Pharmacoeconomics* 2008;26(10):861-77.
225. Riegger GA. Effects of quinapril on exercise tolerance in patients with mild to moderate heart failure. *Eur Heart J* 1991;12(6):705-11.
226. Rodgers A, Chapman N, Woodward M, et al. Perindopril-based blood pressure lowering in individuals with cerebrovascular disease: consistency of benefits by age, sex and region. *J Hypertens* 2004;22:653-59
227. Rodriguez-Granillo GA, Winter S, Bruining N, et al. Effect of perindopril on coronary remodelling: insights from a multicentre, randomized study. *Eur Heart J* 2007;28:2326-31.
228. Rodriguez-Granillo GA, Vos J, Bruining N, et al. Long-term effect of perindopril on coronary atherosclerosis progression (from the perindopril's prospective effect on cornary

Appendix B: List of Excluded Studies

- atherosclerosis by angiography and intravascular ultrasound evaluation (PERSPECTIVE) study. *Am J Cardiol* 2007;100:159-63.
229. Rouleau JL, Chatterjee K, Bengt W, et al. Alterations in left ventricular function and coronary hemodynamics with captopril, hydralazine and prazosin in chronic ischemic heart failure: a comparative study. *Circulation* 1982;65:671-78.
230. Rutherford JD, Pfeffer MA, Moye LA, et al. Effects of captopril on ischemic events after myocardial infarction. Results of the survival and ventricular enlargement trial. SAVE investigators. *Circulation* 1994;90:1731-38.
231. Sabatine MS, Morrow DA, Jablonski KA, et al. Prognostic significance of the centers for disease control/American Heart Association high-sensitivity c-reactive protein cut points for cardiovascular and other outcomes in patients with stable coronary artery disease. *Circulation* 2007;115:1528-36.
232. Sadamatsu K, Shimokawa H, Tashiro H, et al. Long term treatment with enalapril reduces plasma concentrations of macrophage colony stimulating factor in patients with coronary artery disease. *Heart* 2001;86:456-58.
233. Safar ME, Vaisse B, Blacher J, et al. Pulse pressure monitoring of open antihypertensive therapy. *Am J Hypertens* 2004;12(12 Pt 1):1088-94.
234. Schädlich PK, Brecht JG, Rangoonwala B, et al. Cost effectiveness of ramipril in patients at high risk for cardiovascular events: Economic evaluation of the HOPE (Heart Outcomes Prevention Evaluation) study for Germany from the statutory health insurance perspective. *Pharmacoeconomics* 2004;22(25):955-73.
235. Schlaifer JD, Mancini GBJ, O'Neill BJ, et al. Influence of smoking status on angiotensin-converting enzyme inhibitor-related improvement in coronary endothelial function. *Cardiovasc Drugs Ther* 1999;13:201-9.
236. Schlaifer JD, Wargovich TJ, O'Neill B, et al. Effects of quinapril on coronary blood flow in coronary artery disease patients with endothelial dysfunction. *Am J Cardiol* 1997;80(12):1594-7.
237. Schmieder RE. Endothelial dysfunction: how can one intervene at the beginning of the cardiovascular continuum? *J Hypertens* 2006;24(suppl 2):S31-5.
238. Schmieder RE, Handrock R and the VALUE study group. [Control of systolic blood pressure in the VALUE trial after 12 months.] *Dtsch Med Wochenschr* 2003;128(10):485-6. (German).
239. Schrader J, Luders S, Kulschewski A, et al. Morbidity and mortality after stroke, eprosartan compared with nitrendipine for secondary prevention: Principal results of a prospective randomized controlled study (MOSES). *Stroke* 2005;36:1218-26.
240. Schrier RW, Estacio RO. Additional follow-up from the ABCD trial in patients with type 2 diabetes and hypertension. *N Engl J Med* 2000;343(26):1969.
241. Schrier RW, Estacio RO, Esler A, et al. Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. *Kidney Int* 2002;61:1086-97.
242. Schrier RW, Estacio RO, Jeffers B. Appropriate blood pressure control in NIDDM (ABCD) trial. *Diabetologia* 1996;39:1646-54.
243. Scientific Committee of the PERTINENT Sub-Study on behalf of the EUROPA-PERTINENT Investigators. PERTINENT-Perindopril-thrombosis, inflammation, endothelial dysfunction and neurohormonal activation trial: A sub-study of the EUROPA study. *Cardiovasc Drugs Ther* 2003;17:83-91.

Appendix B: List of Excluded Studies

244. Seth S, Mohan A. Quinapril for coronary artery disease. *Natl Med J India* 1997;10(2):73-4.
245. Sever PS, Dahlof B, Poulter NR, et al. Anglo-scandinavian cardiac outcomes trial: a brief history, rationale and outline protocol. *J Hum Hypertens* 2001;15(suppl 1):S11-12
246. Sever PS, Poulter NR, Elliott WJ, et al. Blood pressure reduction is not the only determinant of outcome. *Circulation* 2006;113:2754-74.
247. Simons WR. Comparative cost effectiveness of angiotensin II receptor blockers in a US managed care setting: olmesartan medoxomil compared with losartan, valsartan, and irbesartan. *Pharmacoeconomics* 2003;21(1):61-74.
248. Sleight P. The ONTARGET/TRANSCENT trial programme: baseline data. *Acta Diabetol* 2005;42:S50-6.
249. Smebye ML, Iversen EK, Hoiegggen A, et al. Effect of hemoglobin levels on cardiovascular outcomes in patients with isolated systolic hypertension and left ventricular hypertrophy (from the LIFE study). *Am J Cardiol* 2007;100:855-9.
250. Solomon SD, Anavekar N, Skali H, et al. Influence of ejection fraction on cardiovascular outcomes in a broad spectrum of heart failure patients. *Circulation* 2005;112:3738-44.
251. Solomon SD, Wang D, Finn P, et al. Effect of candesartan on cause-specific mortality in heart failure patients: The candesartan in heart failure assessment of reduction in mortality and morbidity (CHARM) program. *Circulation* 2004;110: 2180-3.
252. Sutton MSJ, Pfeffer MA, Lemuel M, et al. Cardiovascular death and left ventricular remodeling two years after myocardial infarction: Baseline predictors and impact of long-term use of captopril: Information from the survival and ventricular enlargement (SAVE) trial. *Circulation* 1997;96:3294-99.
253. Stoschitzky K. The EUROPA trial. *Lancet* 2003;362:1935-6
254. Stumpe KO, Overlack A, on behalf of the Perindopril Therapeutic Safety Study Group (PUTS). A new trial of the efficacy, tolerability, and safety of angiotensin-converting enzyme inhibition in mild systemic hypertension with concomitant diseases and therapies. *Am J Cardiol* 1993;71:32E-37E.
255. Suzuki H, Kanno Y, Sugahara S, et al. Effect of angiotensin receptor blockers on cardiovascular events in patients undergoing hemodialysis: an open-label randomized controlled trial. *Am J Kidney Dis* 2008;52(3):501-6
256. Swedberg K, Pfeffer M, Granger C, et al. Candesartan in heart failure – Assessment of reduction in mortality and morbidity (CHARM): Rationale and design. *J Card Fail* 1999;5(3):276-82.
257. Tedesco MA, Natale F, Calabro R. Effects of monotherapy and combination therapy on blood pressure control and target organ damage: a randomized prospective intervention study in a large population of hypertensive patients. *J Clin Hypertens* 2006;8(9):634-41.
258. The CAPPP Group. The Captopril Prevention Project: a prospective intervention trial of angiotensin converting enzyme inhibition in the treatment of hypertension. The CAPPP Group. *J Hypertens* 1990;8(11):985-90.
259. The GLANT Study Group. A 12-month comparison of ACE inhibitor and CA antagonist therapy in mild to moderate essential hypertension--The GLANT Study. Study Group on Long-term Antihypertensive Therapy. *Hypertens Res* 1995;18(3):235-44.
260. The Multicenter European Trial with Cilazapril After Angioplasty to Prevent Transluminal Coronary Obstruction and Restenosis (MERCATOR) Study Group. Does the new angiotensin converting enzyme inhibitor Cilazapril Prevent Restenosis After Percutaneous

Appendix B: List of Excluded Studies

- Transluminal Coronary Angioplasty? Results of the MERCATOR Study: A Multicenter, Randomized, Double-Blind Placebo-Controlled Trial. *Circulation* 1992;86(1):100-10.
261. The PHYLLIS Project Group. Plaque Hypertension Lipid-Lowering Italian Study (PHYLLIS): a protocol for non-invasive evaluation of carotid atherosclerosis in hypercholesterolaemic hypertensive subjects. *J Hypertens* 1993;11(suppl 5):S314-5.
 262. The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med* 1992;327(10):685-91
 263. Timmis AD, Pitt B. Effects of ACE inhibitors on coronary atherosclerosis and restenosis. *Br Heart J* 1994;72:57-60.
 264. Toprak O, Cirit M, Bayata S, et al. [The effect of pre-procedural captopril on contrast-induced nephropathy in patients who underwent coronary angiography]. *Anadolu Kardiyol Derg* 2003;3:98-103. (Turkish).
 265. Tousoulis D, Kourtellis P, Antoniadis C, et al. Effects of irbesartan and perindopril on forearm reactive hyperemia and inflammatory process, in normotensive patients with coronary artery disease. *Int J Cardiol* 2008;124:127-9.
 266. Toyofuku M, Imazu M, Sumii K, et al. Influence of angiotensinogen M253T gene polymorphism and an angiotensin converting enzyme inhibitor on restenosis after percutaneous coronary intervention. *Atherosclerosis* 2002;160(2):339-44.
 267. Trenkwalder P, Elmfeldt D, Hofman A, et al. The study on cognition and prognosis in the elderly (SCOPE) – major CV events and stroke in subgroups of patients. *Blood Press* 2005;14(1):31-7.
 268. Trevelyan J, Brull DJ, Needham EW, et al. Effect of enalapril and losartan on cytokines in patients with stable angina pectoris awaiting coronary artery bypass grafting and their interaction with polymorphisms in the interleukin-6 gene. *Am J Cardiol* 2004;94(5):564-69.
 269. Tschope C, Tschope R, Unger T. [Microalbuminuria in patients with hypertension and cardiovascular comorbidity, REAL observational study of treatment with Irbesartan/HCTZ.] *MMW Fortschr Med* 2006;148(25):48-51. (German).
 270. Tzourio C. Vascular factors and cognition: toward a prevention of dementia? *J Hypertens* 2003;21(suppl 5):S15-9.
 271. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998;317:703-13.
 272. Ujiie Y, Hirosaka A, Mitsugi M, et al. Effects of angiotensin-converting enzyme inhibitors or an angiotensin receptor blocker in combination with aspirin and cilostazol on in-stent restenosis. *Int Heart J* 2006;47:173-84.
 273. Umemoto S, Kamiya A, Matsuzaki M, et al. [COPE trial.] *Nippon Rinsho* 2006;64(suppl 6): 470-4. (Japanese).
 274. van den Heuvel ADFM, van Gilst WH, van Veldhuisen DJ, et al. Long-term anti-ischemic effects of angiotensin-converting enzyme inhibition in patients after myocardial infarction. *J Am Coll Cardiol* 1997;30:400-5.
 275. van Haelst PL, Tervaert JW, van Geel P, et al. Long term angiotensin converting enzyme-inhibition in patients after coronary artery bypass grafting reduces levels of soluble intercellular cell adhesion molecule-1. *Eur J Vasc Endovasc Surg* 2003;26:387-91.

Appendix B: List of Excluded Studies

276. Vanuzzo D. [The reduction of cardiovascular risk with ramipril.] *Ital Heart J* 2000;1(suppl 5):713-4. (Italian).
277. Vizir VA, Berezin AE. [Losartan in therapy of chronic heart failure.] *Klin Med* 2000;78(2):36-9. (Russian).
278. Wachtell K, Horneftam B, Lehto M, et al. Cardiovascular morbidity and mortality in hypertensive patients with a history of atrial fibrillation: The losartan intervention for endpoint reduction in hypertension (LIFE) study. *J Am Coll Cardiol* 2005;45(5):705-11.
279. Wang L, Huang X, Zhou Y, et al. [Effects of angiotensin receptor blocker and angiotensin-converting enzyme inhibitor on vascular endothelial function in type 2 diabetic patients with atherosclerosis: a comparative study.] *J First Mil Med Univ* 2005;25:967-71. (Chinese).
280. Weber MA, Julius S, Kjeldsen SE, et al. Blood pressure dependent and independent effects of antihypertensive treatment on clinical events in the VALUE trial. *Lancet* 2004;363:2049-51.
281. Whelton PK, Barzilay J, Cushman WC, et al. Clinical outcomes in antihypertensive treatment of type 2 diabetes, impaired fasting glucose concentration, and normoglycemia. *Arch Intern Med* 2005;165:1401-9.
282. White HD, Aylward PEG, Huang Z, et al. Mortality and morbidity remain high despite captopril and/or valsartan therapy in elderly patients with left ventricular systolic dysfunction, heart failure, or both after acute myocardial infarction: Results from the valsartan in acute myocardial infarction trial (VALIANT). *Circulation* 2005;112:3391-9.
283. Wiedermann CJ. Losartan for cardiovascular disease in patients with and without diabetes in the LIFE study. *Lancet* 2002;359:2199.
284. Wierchowieski M, Filipiak J, Poprawski K, et al. [The effect of on-year Captopril treatment on exercise tolerance and myocardial ischemia in patients with myocardial infarction.] *Pol Merk Lek* 2000;9:536-40. (Polish).
285. Willenheimer R, Juul-Moller S, Forslund L, et al. No effects on myocardial ischaemia in patients with stable ischaemic heart disease after treatment with ramipril for 6 months. *Curr Control Trials Cardiovasc Med* 2001;2:99-105.
286. Willenheimer R, Rydberg E, Oberg L, et al. ACE inhibition with ramipril improves left ventricular function at rest and post exercise in patients with stable ischemic heart disease and preserved left ventricular systolic function. *Eur Heart J* 1999;20:1647-56.
287. Wing LMH, Reid CM, Ryan P, et al. A comparison of outcomes with angiotensin-converting-enzyme inhibitors and diuretics for hypertension in the elderly. *N Engl J Med* 2003;348:583-92 .
288. Wing LM, Reid CM, Ryan P, et al. Second Australian National Blood Pressure Study (ANBP2). Australian Comparative Outcome Trial of ACE inhibitor- and diuretic-based treatment of hypertension in the elderly. Management Committee on behalf of the High Blood Pressure Research Council of Australia. *Clin and Exper Hypertension* 1997;19(5-6):779-91.
289. Wing LM, Reid CM, Ryan P, et al. Second Australian National Blood Pressure Study (ANBP2). Australian Comparative Outcome Trial of ACE inhibitor- and diuretic-based treatment of hypertension in the elderly. Management Committee on behalf of the High Blood Pressure Research Council of Australia. *Clin and Exper Hypertension* 1997;19(5-6):779-91.

Appendix B: List of Excluded Studies

290. Wright JT, Dunn JK, Cutler JA, et al. Outcomes in hypertensive black and nonblack patients treated with chlorthalidone, amlodipine, and lisinopril. *JAMA* 2005;293:1595-608.
291. Xie Z, Zeng S. [Changes of plasma soluble CD40 ligand, tumor necrosis factor alpha and C reactive protein in patients with coronary heart disease and the intervention of losartan.] *Zhongguo Linchuang Kangfu* 2005;9:71-3 (Chinese).
292. Yamabe T, Imazu M, Yamamoto H, et al. Effects of cilazapril on vascular restenosis after percutaneous transluminal coronary angioplasty. *Coron Artery Dis* 1995;6(7):573-9.
293. Yip GWK, Wang M, Wang T, et al. The Hong Kong diastolic heart failure study: A randomised controlled trial of diuretics, irbesartan and ramipril on quality of life, exercise capacity, left ventricular global and regional function in heart failure with a normal ejection fraction. *Heart* 2008;94:573-80.
294. Yoshida D, Higashiura K, Shimamoto K. [Treatment of hypertensive patients with impaired glucose tolerance (IGT).] *Nippon Rinsho* 2005;63(suppl 2):533-7. (Japanese).
295. Young JB. ACE inhibitors and ischemic heart disease. *Coron Artery Dis* 1995;6(4):272-80.
296. Yusuf S, Gerstein H, Hoogwerf B, et al. Ramipril and the development of diabetes. *JAMA* 2001;286:1882-5.
297. Yusuf S, Pfeffer MA, Swedberg K, et al. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved trial. *Lancet* 2003;362:777-81.
298. Yusuf S, Diener HC, Sacco RL, et al for the PROfESS Study Group. Telmisartan to prevent recurrent stroke and cardiovascular events. *N Engl J Med* 2008;359(12):1225-37.
299. Zamorski MA. CCBs vs ACE inhibitors in patients with diabetes. *J Fam Pract* 1998;47:12-3.

Appendix B: List of Excluded Studies

Systematic Review Search

1. Abdulla J, Barlera S, Latini R, et al. A systematic review: effect of angiotensin converting enzyme inhibition on left ventricular volumes and ejection fraction in patients with a myocardial infarction and in patients with left ventricular dysfunction. *Eur J Heart Failure*. 2007;9(2):129-35.
2. Abuissa H, Jones PG, Marso SP, et al. Angiotensin-converting enzyme inhibitors or angiotensin receptor blockers for prevention of type 2 diabetes: a meta-analysis of randomized clinical trials. *J Am Coll Cardiol*. 2005;46(5):821-6.
3. Aguilar MI, Hart R, Pearce LA. Oral anticoagulants versus antiplatelet therapy for preventing stroke in patients with non-valvular atrial fibrillation and no history of stroke or transient ischemic attacks. *Cochrane Database of Systematic Reviews*. 4, 2008.
4. Amsallem E, Kasparian C, Haddour G, et al. Phosphodiesterase III inhibitors in heart failure. *Cochrane Database of Systematic Reviews*. 4, 2008.
5. Anarow WS. Treatment of congestive heart failure in older persons. *J Am Geriatr Soc*. 1997;45(10):1252-7.
6. Andraws R, Brown DL. Effect of inhibition of the renin-angiotensin system on development of type 2 diabetes mellitus (meta-analysis of randomized trials). *Am J Cardiol*. 2007;99(7):1006-12.
7. Arici M, Erdem Y. Dual blockade of the renin-angiotensin system for cardiorenal protection: an update. *Am J Kidney Dis* 2009;53(2):332-45.
8. Aronow WS. Management of older persons after myocardial infarction. *J Am Geriatr Soc*. 1998;46(11):1459-68.
9. Aronow WS. Treatment of congestive heart failure in older persons. *J Am Geriatr Soc*. 1997;47(10):1252-7.
10. Aronson D, Rayfield EJ, Chesebro JH. Mechanisms determining course and outcome of diabetic patients who have had acute myocardial infarction. *Ann Intern Med*. 1997;126(4):296-306.
11. Aursnes I, Tvette IF, Gasemyr J, et al. Clinical efficacies of antihypertensive drugs. *Scand Cardiovasc J*. 2003;37(2):72-9.
12. Baillie GM, Sherer JT, Weart CW. Insulin and coronary artery disease: is syndrome X the unifying hypothesis? *Ann Pharmacother*. 1998;32(2):233-47.
13. Baker DW. Prevention of heart failure. *J Cardiac Failure*. 2002;8(5):333-46.
14. Berry C, Tardif JC, Bourassa MG. Coronary heart disease in patients with diabetes: part I: recent advances in prevention and noninvasive management. *J Am Coll Cardiol*. 2007;49(6):631-42.
15. Bethel MA, Holman R, Haffner SM, et al. Determining the most appropriate components for a composite clinical trial outcome. *Am Heart J*. 2008;56(4):633-40.
16. Bjelakovic G, Nikolova D, Gluud LL, et al. Antioxidant supplements for preventing gastrointestinal cancers. *Cochrane Database of Systematic Reviews*. 4, 2008.
17. Bjelakovic G, Nikolova D, Gluud LL, et al. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database of Systematic Reviews*. 4, 2008.

Appendix B: List of Excluded Studies

18. Blood Pressure Lowering Treatment Trialists' Collaboration. Blood pressure-dependent and independent effects of agents that inhibit the renin-angiotensin system. *J Hypertension*. 2007;25(5):951-8.
19. Boden WE, Vray M, Eschwege E, et al. Heart rate-lowering and -regulating effects of once-daily sustained-release diltiazem. *Clin Cardiol*. 2001;24(1):73-9.
20. Boos CJ. Cardiovascular protection with ace inhibitors--more HOPE for EUROPA? *Med Sci Monitor*. 2004;10(12):SR23-8.
21. Bugiardini R, Bairey Merz CN. Angina with "normal" coronary arteries: a changing philosophy. *JAMA*. 2005;293(4):477-84.
22. Butler R, Morris AD, Struthers AD. Angiotensin-converting enzyme gene polymorphism and cardiovascular disease. *Clin Sci*. 1997;93(5):391-400
23. Chen H, Liu J, Yang M. Corticosteroids for viral myocarditis. *Cochrane Database of Systematic Reviews*. 4, 2008.
24. Chen J, Wu G, Li S, et al. Shengmai (a traditional Chinese herbal medicine) for heart failure. *Cochrane Database of Systematic Reviews*. 4, 2008.
25. Coleman CI, Baker WL, Kluger J, et al. Antihypertensive medication and their impact on cancer incidence: a mixed treatment comparison meta-analysis of randomized controlled trials. *J Hypertension*. 2008;26(4):622-9.
26. Dahlof B, Devereux RB, Kjeldsen SE, et al. Atenolol as a comparator in outcome trials in hypertension: a correct choice in the past, but not for the future? *Blood Pressure*. 2007;16(1):6-12.
27. de Courten B, Barber MN, Johnston RV, et al. Hypolipidemic and antihypertensive drugs for prevention of cardiovascular complications in patients with rheumatoid arthritis. *Cochrane Database of Systematic Reviews*. 4, 2008.
28. de Vries RJ, van Veldhuisen DJ, Dunselman PH. Efficacy and safety of calcium channel blockers in heart failure: focus on recent trials with second-generation dihydropyridines. *Am Heart J*. 2000;139(2):185-94.
29. Dicipinigaitis PV. Angiotensin-converting enzyme inhibitor-induced cough: ACCP evidence-based clinical practice guidelines. *Chest*. 2006;129(1 Suppl):169S-173S.
30. Dobre D, van Veldhuisen DJ, DeJongste MJ, et al. The contribution of observational studies to the knowledge of drug effectiveness in heart failure. *Br J Clin Pharmacol*. 2007;64(4):406-14.
31. Dornbrook-Lavender KA, Roth MT, Pieper JA. Secondary prevention of coronary heart disease in the elderly. *Ann Pharmacother*. 2003;37(12):1867-76.
32. Dzau VJ. Angiotensin converting enzyme inhibitors and the cardiovascular system. *J Hypertension*. 1992;10(3):S3-10.
33. Elsayed EF, Tighiouart H, Griffith J, et al. Cardiovascular disease and subsequent kidney disease. *Arch Intern Med*. 2007;167(11):1130-6.
34. Faris R, Flather MD, Purcell H, et al. Diuretics for heart failure. *Cochrane Database of Systematic Reviews*. 4, 2008.
35. Filion KB, Pilote L, Rahme E, et al. Perioperative use of cardiac medical therapy among patients undergoing coronary artery bypass graft surgery: a systematic review. *Am Heart J*. 2007;154(3):407-14.
36. Fretheim A, Bjorndal A, Oxman AD, et al. [Which antihypertensive drugs should be used in the primary prevention of cardiovascular disease?] *Tidsskrift for Den Norske Laegeforening*. 2002;122(23):2283-6.

Appendix B: List of Excluded Studies

37. Gillespie EL, White CM, Kardas M, et al. The impact of ACE inhibitors or angiotensin II type 1 receptor blockers on the development of new-onset type 2 diabetes. *Diabetes Care*. 2005;28(9):2261-6.
38. Gluckman TJ, Sachdev M, Schulman SP, et al. A simplified approach to the management of non-ST-segment elevation acute coronary syndromes. *JAMA*. 2005;293(3):349-57.
39. Goderis G, Boland B. Cardiovascular prevention in type 2 diabetic patients: review of efficacious treatments. *Acta Clinica Belgica*. 2004;59(6):329-39.
40. Hankey GJ, Norman PE, Eikelboom JW. Medical treatment of peripheral arterial disease. *JAMA*. 2006;295(5):547-53.
41. Haynes RB, Ackloo E, Sahota N, et al. Interventions for enhancing medication adherence. *Cochrane Database of Systematic Reviews*. 4, 2008.
42. Healey JS, Baranchuk A, Crystal E, et al. Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis. *J Am Coll Cardiol*. 2005;45(11):1832-9.
43. Hilleman DE, Banakar UV. Issues in contemporary drug delivery. Part VI: Advanced cardiac drug formulations. *J Pharm Tech*. 1992;8(5):203-11.
44. Hood WB, Dans AL, Guyatt GH, et al. Digitalis for treatment of congesting heart failure in patients in sinus rhythm. *Cochrane Database of Systematic Reviews*. 4, 2008.
45. Howes LG, Lykos D, Rennie GC. Effects of antihypertensive drugs on coronary artery disease risk: a meta-analysis. *Clin Exp Pharmacol Physiol*. 1996;23(6-7):555-8.
46. Johnson AD, Gong Y, Wang D, et al. Promoter polymorphisms in ACE (angiotensin I-converting enzyme) associated with clinical outcomes in hypertension. *Clin Pharmacol Ther* 2009;85(1):36-44.
47. Kalus JS, White CM. Amlodipine versus Angiotensin-receptor blockers for nonhypertension indications. *Ann Pharmacother*. 2002;36(11):1759-66
48. Kerr CP. Hypertension in the 1990s: a new disease. *J Am Board Fam Prac*. 1993;6(3):243-54.
49. Kikano GE, Brown MT. Antiplatelet therapy for atherothrombotic disease: an update for the primary care physician. *Mayo Clinic Proc*. 2007;82(5):583-93.
50. Kramer JM, Newby LK, Chang WC, et al. International variation in the use of evidence-based medicines for acute coronary syndromes. *Eur Heart J*. 2003;24(23):2133-41.
51. Kuchenbecker R, Berwanger O, Rosito G. Statins for preventing major vascular events in people with hypertension. *Cochrane Database of Systematic Reviews*. 4, 2008.
52. Lafuente-Lafuente C, Mouly S, Longas-Tejero MA, et al. Antiarrhythmics for maintaining sinus rhythm after cardioversion of atrial fibrillation. *Cochrane Database of Systematic Reviews*. 4, 2008.
53. Li J, Zhang Q, Zhang M, et al. Intravenous magnesium for acute myocardial infarction. *Cochrane Database of Systematic Reviews*. 4, 2008.
54. Lip GY, Makin AJ. Treatment of hypertension in peripheral arterial disease. *Cochrane Database of Systematic Reviews*. 4, 2008.
55. Lip GYH, Chung I. Anticoagulation for heart failure in sinus rhythm. *Cochrane Database of Systematic Reviews*. 4, 2008.
56. Lip GYH, Chung I. Antiplatelet agents versus control or anticoagulation for heart failure in sinus rhythm. *Cochrane Database of Systematic Reviews*. 4, 2008.
57. Liu G, Chen X, Qu T. Huangqi preparations for unstable angina. *Cochrane Database of Systematic Reviews*. 4, 2008.

Appendix B: List of Excluded Studies

58. Liu X, Li J, Wu T. Nitrates for stable angina. *Cochrane Database of Systematic Reviews*. 4, 2008.
59. Loomba RS, Arora R. Prevention of coronary heart disease in women. *Ther Adv Cardiovasc Dis* 2008;2(5):321-7.
60. Madrid AH, Peng J, Zamora J, et al. The role of angiotensin receptor blockers and/or angiotensin converting enzyme inhibitors in the prevention of atrial fibrillation in patients with cardiovascular diseases: meta-analysis of randomized controlled clinical trials. *Pacing and Clin Electrophys*. 2004;27(10):1405-10.
61. Mancini GB. Emerging role of angiotensin II type 1 receptor blockers for the treatment of endothelial dysfunction and vascular inflammation. *Can J Cardiol*. 2002;18(12):1309-16.
62. Mancini GB. Long-term use of angiotensin-converting enzyme inhibitors to modify endothelial dysfunction: a review of clinical investigations. *Clin Invest Med*. 2000;23(2):144-61.
63. Mark L, Dani G, Kiss Z, et al. [A change of attitude in lipidology, achievement of target levels. What comes next?][Hungarian]. *Orvosi Hetilap* 2008;149(37):1731-6.
64. Matchar DB, McCrory DC, Orlando LA, et al. Systematic review: comparative effectiveness of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers for treating essential hypertension. *Ann Intern Med*. 2008;148(1):16-29.
65. McCall KL, Craddock D, Edwards K. Effect of angiotensin-converting enzyme inhibitors and angiotensin II type 1 receptor blockers on the rate of new-onset diabetes mellitus: a review and pooled analysis. *Pharmacotherapy*. 2006;26(9):1297-306.
66. McDowell SE, Coleman JJ, Ferner RE. Systematic review and meta-analysis of ethnic differences in risks of adverse reactions to drugs used in cardiovascular medicine. *BMJ*. 2006;332(7551):1177-81.
67. McGuinness B, Todd S, Passmore P, et al. Blood pressure lowering in patients without prior cerebrovascular disease for prevention of cognitive impairment and dementia. *Cochrane Database of Systematic Reviews*. 4, 2008.
68. Messerli FH, Grossman E, Goldbourt U. Antihypertensive therapy in diabetic hypertensive patients. *Am J Hypertension*. 2001;14(5):S12-16.
69. Meune C, Mahe I, Mourad JJ, et al. Interaction between angiotensin-converting enzyme inhibitors and aspirin: a review. *Eur J Clin Pharmacol*. 2000;56(9-10):609-20.
70. Mulrow CD, Mulrow JP, Linn WD, et al. Relative efficacy of vasodilator therapy in chronic congestive heart failure. Implications of randomized trials. *JAMA*. 1988;259(23):3422-6.
71. Nawarskas KK, Spinler SA. Update on the interaction between aspirin and angiotensin-converting enzyme inhibitors. *Pharmacotherapy*. 2000;20(6):698-710.
72. Neal B, MacMahon S, Chapman N, et al. Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Lancet* 2000;356(9246):1955-64.
73. Nelson KM, Yeager BF. What is the role of angiotensin-converting enzyme inhibitors in congestive heart failure and after myocardial infarction? *Ann Pharmacother*. 1996;30(9):986-93.
74. Nicolai SPA, Kruidenier LM, Bendermacher BLW, et al. Gingko biloba for intermittent claudication. *Cochrane Database of Systematic Reviews*. 4, 2008.

Appendix B: List of Excluded Studies

75. Nolan RP, Jong P, Barry-Bianchi SM, et al. Biobehavioral and exercise therapies on heart rate variability in coronary artery disease: a systematic review. *Eur J Cardiovasc Prevent Rehab.* 2008;15(4):386-96.
76. Nony P, Boissel JP, Girard P, et al. Relative efficacy of angiotensin converting enzyme inhibitors on mortality of patients with congestive heart failure: implications of randomized trials and role of the aetiology (ischaemic or non-ischaemic) of heart failure. *Eur Heart J.* 1992;13(8):1101-8.
77. Okrainec K, Platt R, Pilote L, et al. Cardiac medical therapy in patients after undergoing coronary artery bypass graft surgery: a review of randomized controlled trials. *J Am Coll Cardiol.* 2005;45(2):177-84.
78. Pahor M, Psaty BM, Alderman MH, et al. Health outcomes associated with calcium antagonists compared with other first-line antihypertensive therapies: a meta-analysis of randomised controlled trials. *Lancet.* 2000;356(9246):1949-54.
79. Pahor M, Psaty BM, Alderman MH, et al. Therapeutic benefits of ACE inhibitors and other antihypertensive drugs in patients with type 2 diabetes. *Diabetes Care.* 2000;23(7):888-92.
80. Paramothayan NS, Lasserson TJ, Wells AU, et al. Prostacyclin for pulmonary hypertension in adults. *Cochrane Database of Systematic Reviews.* 4, 2008.
81. Paravastu S, Chandra V, Mendonca D, et al. Beta blockers for peripheral arterial disease. *Cochrane Database of Systematic Reviews.* 4, 2008.
82. Park IU, Taylor AL. Race and ethnicity in trials of antihypertensive therapy to prevent cardiovascular outcomes: a systematic review. *Ann Fam Med.* 2007;5(5):444-52.
83. Pedersen SA, Galatius S, Olsen MH, et al. High prevalence of risk factors in coronary artery disease in EUROPA gives HOPE for ACE inhibitors after PEACE. *Cardiology.* 2008;111(1):63-7.
84. Perez MI, Musini VM, Wright JM. Effect of early treatment with anti-hypertensive drugs on short and long-term health outcomes in patients with an acute cardiovascular event. *Cochrane Database of Systematic Reviews.* 4, 2008.
85. Perez MI, Musini VM, Wright JM. Pharmacological interventions for hypertensive emergencies. *Cochrane Database of Systematic Reviews.* 4, 2008.
86. Pillay A, O'Reagan LL. Methyldopa for the management of essential hypertension. *Cochrane Database of Systematic Reviews.* 4, 2008.
87. Psaty BM, Lumley T, Furberg CD, et al. Health outcomes associated with various antihypertensive therapies used as first-line agents: a network meta-analysis. *JAMA.* 2003;289(19):2534-44.
88. Psaty BM, Smith NL, Siscovick DS, et al. Health outcomes associated with antihypertensive therapies used as first-line agents. A systematic review and meta-analysis. *JAMA.* 1997;277(9):739-45
89. Reboldi G, Angeli F, Cavallini C, et al. Comparison between angiotensin-converting enzyme inhibitors and angiotensin receptor blockers on the risk of myocardial infarction, stroke and death: a meta-analysis. *J Hypertension.* 2008;26(7):1282-9.
90. Ribeiro AB. Angiotensin II antagonists--therapeutic benefits spanning the cardiovascular disease continuum from hypertension to heart failure and diabetic nephropathy. *Curr Med Res Opin.* 2006;22(1):1-16.
91. Richter B, Bandeira-Echtler E, Bergerhoff K, et al. Pioglitazone for type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews.* 4, 2008.

Appendix B: List of Excluded Studies

92. Richter B, Bandeira-Echtler E, Bergerhoff K, et al. Dipeptidyl peptidase-4 (DPP-4) inhibitors for type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*. 4, 2008.
93. Richter B, Bandeira-Echtler E, Bergerhoff K, et al. Rosiglitazone for type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*. 4, 2008.
94. Ruilope LM, Segura J. Advantages of new cardiovascular risk-assessment strategies in high-risk patients with hypertension. *Clin Ther*. 2005;27(10):1658-68.
95. Scharplatz M, Puhan MA, Steurer J, et al. What is the impact of the ACE gene insertion/deletion (I/D) polymorphism on the clinical effectiveness and adverse events of ACE inhibitors?--Protocol of a systematic review. *BCM Medical Genetics*. 2004;5:23.
96. Shu D, Dong B, Wu T, et al. Beta blockers for stable angina. *Cochrane Database of Systematic Reviews*. 4, 2008.
97. Siragy H, Huang J, Lieb DC. The development of the direct renin inhibitor aliskiren: treating hypertension and beyond. *Exp Opin Emerg Drugs* 2008;13(3):417-30.
98. Sleight P, Jakobsen A, Heroys J, et al. No HOPE without proof: do ARBs meet the standard for cardiovascular protection? *Medscape J Med* 2008;10 Suppl:s6.
99. Souza LM, Atallah An, Saconato H, et al. Oral drugs for hypertensive urgencies. *Cochrane Database of Systematic Reviews*. 4, 2008.
100. Staessen JA, Wang JG, Thijs L. Cardiovascular protection and blood pressure reduction: a meta-analysis. *Lancet*. 2001;358(9290):1305-15.
101. Strippoli GFM, Bonifati C, Craig M, et al. Angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists for preventing the progression on diabetic kidney disease. *Cochrane Database of Systematic Reviews*. 4, 2008.
102. Strippoli GFM, Craig M, Craig JC. Antihypertensive agents for preventing diabetic kidney disease. *Cochrane Database of Systematic Reviews*. 4, 2008.
103. Strippoli GFM, Craig M, Schena FP, et al. Antihypertensive agents for preventing progression of diabetic kidney disease. *Cochrane Database of Systematic Reviews*. 4, 2008.
104. Thorp ML, Ditmer DG, Nash MK, et al. A study of the prevalence of significant increases in serum creatinine following angiotension converting enzyme inhibitor administration. *J Human Hypertension*. 2005;19(5):389-92.
105. Turnbull F, Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomised trials. *Lancet*. 2003;362(9395):1527-35.
106. Turnbull F, Neal B, Algert C, et al. Effects of different blood pressure-lowering regimens on major cardiovascular events in individuals with and without diabetes mellitus: results of prospectively designed overviews of randomized trials. *Arch Intern Med*. 2005;165(12):1410-9.
107. Verdecchia P, Reboldi G, Angeli F, et al. Angiotensin-converting enzyme inhibitors and calcium channel blockers for coronary heart disease and stroke prevention. *Hypertension*. 2005;46(2):386-92.
108. Vinjar B, Stewart M. Oral vasodilators for primary Raynaud's phenomenon. *Cochrane Database of Systematic Reviews*. 4, 2008.
109. Vuong AD, Annis LG. Ramipril for the prevention and treatment of cardiovascular disease. *Ann Pharmacother*. 2003;37(37):412-9.
110. Wang JG, Li Y, Franklin SS, et al. Prevention of stroke and myocardial infarction by amlodipine and Angiotensin receptor blockers: a quantitative overview. *Hypertension*. 2007;50(1):181-8.

Appendix B: List of Excluded Studies

111. Wang JG, Staessen JA, Li Y, et al. Carotid intima-media thickness and antihypertensive treatment: a meta-analysis of randomized controlled trials. *Stroke*. 2006;37(7):1933-40.
112. Wang JG, Staessen JA. Benefits of antihypertensive pharmacologic therapy and blood pressure reduction in outcome trials. *J Clin Hypertension*. 2003;5(1):66-75.
113. Weir MR. Effects of renin-angiotensin system inhibition on end-organ protection: can we do better? *Clin Therapeutics*. 2007;29(9):1803-24.
114. Wilson MD, Weart CW. Hypertension: are beta-blockers and diuretics appropriate first-line therapies? *Ann Pharmacother*. 1994;28(5):617-25.
115. Wiysonge C, Shey U, Bradley HA, et al. Beta-blockers for hypertension. *Cochrane Database of Systematic Reviews*. 4, 2008.
116. Wu T, Harrison RA, Chen X, et al. Tongxinluo (Tong xin Luo or Tong-xin-luo) capsule for unstable angina pectoris. *Cochrane Database of Systematic Reviews*. 4, 2008.
117. Zannad F, Jakobsen A, Heroys J, et al. Cardiovascular high-risk patients – treat to protect, but whom? *Medscape J Med* 2008;10 Suppl:s2.
118. Zheng J, Li J, Liu GJ, et al. Traditional Chinese interventions for stable angina. *Cochrane Database of Systematic Reviews*. 4, 2008

Appendix C: Additional Evidence Tables and Analyses

Abbreviations

| Acronym/Abbreviation | Definition |
|----------------------|---|
| ACE | Angiotensin Converting Enzyme |
| ACEI | Angiotensin Converting Enzyme Inhibitor |
| ADE | Adverse Drug Event |
| AHR | Adjusted Hazard Ratio |
| AMSTAR | Assess the Methodological quality of SysteMatic Review |
| APRES | Angiotensin-converting Enzyme inhibition Post Revascularization Study |
| ARB | Angiotensin Receptor Blocker |
| CABG | Coronary Artery Bypass Grafting |
| CAD | Coronary Artery Disease |
| CAMELOT | Comparison of Amlodipine vs Enalapril to Limit Occurrences of Thrombosis |
| CCB | Calcium Channel Blocker |
| CHF | Congestive Heart Failure |
| CI | Confidence Interval |
| CV | Cardiovascular |
| DM | Diabetes Mellitus |
| EKG | Electrocardiogram |
| EUROPA | EUROpean trial On reduction of cardiac events with Perindopril in stable coronary Artery disease |
| FOSIDIAL | FOSInopril in DIALysis |
| F/U | Follow-Up |
| GRADE | Grading of Recommendations Assessment, DEvelopment |
| HF | Heart Failure |
| HOPE | Heart Outcomes Prevention Evaluation |
| HR | Hazard Ratio |
| HTN | Hypertension |
| IC | Intermittent Claudication |
| IHD | Ischemic Heart Disease |
| IMAGINE | Ischemia Management with Accupril post-bypass Graft via Inhibition of the coNverting Enzyme |
| JMIC-B | Japan Multicenter Investigation for Cardiovascular Diseases-B |
| LVEF | Left Ventricular Ejection Fraction |
| LVH | Left Ventricular Hypertrophy |
| MARCATOR | Multicenter American Research trial with Cilazapril After angioplasty to prevent Transluminal coronary Obstruction and Restenosis |
| MI | Myocardial Infarction |
| N/A | Not Applicable |
| NR | Not Reported |
| ONTARGET | ONgoing Telmisartan Alone in combination with Ramipril Global Endpoint Trial |
| OR | Odds Ratio |
| PARIS | Effect of ACE inhibitors on angiographic restenosis after coronary stenting |
| PART-2 | Prevention of Atherosclerosis with Ramipril Trial-2 |
| PCI | Percutaneous Coronary Intervention |

Appendix C: Additional Evidence Tables and Analyses

| | |
|----------------|--|
| PEACE | Prevention of Events with Angiotensin Converting Enzyme Inhibition |
| PTCA | Percutaneous Transluminal Coronary Angioplasty |
| PVD | Peripheral Vascular Disease |
| QUIET | Quinapril Ischemic Event Trial |
| RCT | Randomized Controlled Trial |
| RR | Relative Risk |
| SB | Single Blind |
| SCAT | Simvastatin/enalapril Coronary Atherosclerosis Trial |
| SMILE-ISCHEMIA | Survival of Myocardial Infarction Long-term Evaluation-ISCHEMIA |
| SMT | Standard Medical Therapy |
| TIA | Transient Ischemic Attack |
| TRANSCEND | Telmisartan Randomized Assessment Study in ACE Intolerant subjects with cardiovascular Disease |
| SCR | Scientific Resource Center |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 1. Pertinent Systematic Reviews**

| Reference | Inclusion Criteria | Total Studies Included | Total Pts Included | AMSTAR rating |
|------------------------------|--|-------------------------------|---------------------------|----------------------|
| Al-Mallah 2006 ⁹² | All randomized, placebo controlled trials of ACEIs use in patients with CAD and preserved LV function (LVEF \geq 40%) | 6 | 33,500 | 7/11 |
| Dagenais 2006 ⁹³ | HOPE, EUROPA and PEACE (the three main large trials of ACEIs in patients with atherosclerosis, but without heart failure or LSVD) | 3 | 29,805 | 2/11 |
| Danchin 2006 ⁹⁴ | All placebo-controlled randomized trials with a follow-up of 2 years or longer performed in patients who had stable CAD and either no signs or symptoms of heart failure or no documented LV dysfunction (no LVEF $<$ 35%) | 7 | 33,960 | 9/11 |
| Saha 2007 ⁹⁵ | All randomized, placebo controlled clinical trials with mean study duration of at least 12 months, a use of a tissue-selective ACEI (ramipril, perindopril, quinapril, or trandolapril), and strict inclusion of patients with cardiovascular disease who either had documented EKG evidence of normal left ventricular function (LVEF $>$ 40%) or had no clinical symptoms of CHF at the time of randomization | 4 | 31,555 | 7/11 |
| Lang 2008 ⁹⁶ | All randomized, placebo controlled clinical trials with mean study duration of at least 12 months, a use of a tissue-selective ACEI (ramipril, perindopril, quinapril, or trandolapril), patients with documented DM with evidence of normal left ventricular systolic function or who had no symptoms of congestive heart failure at the onset of the study, and risk factors in addition to DM, according to the Framingham classification | 4 | 10,328 | 6/11 |
| Saha 2008 ⁹⁷ | All RCTs with mean follow-up period of at least 12 months, and that compared effects of tissue-selective ACEI (ramipril, perindopril, quinapril, or trandolapril), with placebo, in patients with known DM who either had documented evidence of normal left ventricular systolic function or had no clinical symptoms of congestive heart failure at the start of the study | 4 | 10,328 | 6/11 |

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 2. KQ1 Total Mortality - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--|------------------------------------|-------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 CV Risk Factor | Death from any cause | Ramipril Placebo | 482/4645 569/4652 | RR 0.84 (0.75 to 0.95) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | All clinical events resulting in death | Ramipril Placebo | 16/308 25/309 | RR 0.64 (0.34 to 1.20) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Death | Enalapril Placebo | 8/229 11/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | Total mortality | Perindopril Placebo | 375/6110 420/6108 | 1-RR 11% (-2% to 23%) |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Cardiovascular + non-cardiovascular deaths | Candesartan Control | 4/194 11/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | All-cause mortality | Enalapril Amlodipine Placebo | 8/673 7/663 6/655 | HR 1.26 (0.44 to 3.65) [†] HR 0.92 (0.33 to 2.53) [¶] HR 1.14 (0.38 to 3.40) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | Total mortality | ACEI [∞] Nifedipine | 15/822 12/828 | RR 0.76 (0.35 to 1.63) [∂] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Death from any cause | Trandolapril Placebo | 299/4158 334/4132 | HR 0.89 (0.76 to 1.04) |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | All cause death | Fosinopril Placebo | 53/196 50/201 | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | Mortality | Candesartan Control | 0/43 7/37 | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Total mortality | Telmisartan Placebo | 364/2954 349/2972 | AHR 1.05 (0.91 to 1.22) |

† = Enalapril vs placebo; ¶ = Amlodipine vs enalapril; ‡ = Amlodipine vs placebo; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; * = Clinical outcome data provided by FOSIDIAL corresponding author; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 3. KQ1 Cardiovascular Mortality - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|-----------------------------------|------------------------------------|-------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 CV Risk Factor | Death from cardiovascular causes | Ramipril Placebo | 282/4645 377/4652 | RR 0.74 (0.64 to 0.87) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Death from cardiovascular disease | Ramipril Placebo | 8/308 18/309 | RR 0.45 (0.19 to 1.03) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Cardiac death | Enalapril Placebo | 4/229 7/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | Cardiovascular mortality | Perindopril Placebo | 215/6110 249/6108 | 1-RR 14% (-3 to 28) |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Cardiovascular death | Candesartan Control | 2/194 9/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Cardiovascular death | Enalapril Amlodipine Placebo | 5/673 5/663 2/655 | HR 2.33 (0.45 to 12.1) [†] HR 1.07 (0.31 to 3.70) [¶] HR 2.46 (0.48 to 12.7) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | Cardiac death or sudden death | ACEI [∞] Nifedipine | 6/822 6/828 | RR 0.96 (0.31 to 3.04) [∂] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Death from cardiovascular causes | Trandolapril Placebo | 146/4158 152/4132 | HR 0.95 (0.76 to 1.19) |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | Cardiovascular death | Fosinopril Placebo | 32/196 31/201 | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Cardiovascular death | Telmisartan Placebo | 227/2954 223/2972 | AHR 1.03 (0.85 to 1.24) |

† = Enalapril vs placebo; ¶ = Amlodipine vs enalapril; ‡ = Amlodipine vs placebo; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; * = Clinical outcome data provided by FOSIDIAL corresponding author; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 4. KQ1 Nonfatal Myocardial Infarction - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|---|------------------------------------|----------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 CV Risk Factor | Acute MI not resulting in death | Ramipril Placebo | 260/4645 333/4652 | 1-RR 23% (9 to 34) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Nonfatal MI requiring hospital admission | Ramipril Placebo | 18/308 19/309 | RR 0.94 (0.49 to 1.80) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Nonfatal MI | Enalapril Placebo | 7/229 12/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) w/o HF | Nonfatal MI (see total MI for definition) | Perindopril Placebo | 295/6110 378/6108 | 1-RR 22% (10 to 33) |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Nonfatal MI | Candesartan Control | 2/194 1/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Nonfatal MI | Enalapril Amlodipine Placebo | 11/673 14/663 19/655 | HR 0.55 (0.26 to 1.15) [†] HR 1.32 (0.60 to 2.90) [¶] HR 0.73 (0.37 to 1.46) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Nonfatal MI | Trandolapril Placebo | 222/4158 220/4132 | HR 1.00 (0.83 to 1.20) |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | Nonfatal MI | Fosinopril Placebo | 9/196 7/201 | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, CV disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

† = Enalapril vs placebo; ‡ = Amlodipine vs placebo; ¶ = Amlodipine vs enalapril; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; * = Clinical outcome data provided by FOSIDIAL corresponding author

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 5. KQ1 Stroke - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--|------------------------------------|--------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Stroke | Ramipril Placebo | 156/4645 226/4652 | RR 0.68 (0.56 to 0.84) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Nonfatal stroke requiring hospital admission | Ramipril Placebo | 7/308 4/309 | RR 1.67 (0.48 to 5.75) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, ↑ cholesterol | Stroke | Enalapril Placebo | 2/229 9/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) w/o HF | Stroke | Perindopril Placebo | 98/6110 102/6108 | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Stroke or TIA | Enalapril Amlodipine Placebo | 8/673 6/663 12/655 | HR 0.66 (0.27 to 1.62) [†] HR 0.76 (0.26 to 2.20) [¶] HR 0.50 (0.19 to 1.32) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | HTN and CAD | Cerebrovascular accidents | ACEI [∞] Nifedipine | 16/822 16/828 | RR 0.76 (0.56 to 2.02) [∂] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Stroke | Trandolapril Placebo | 71/4158 92/4132 | HR 0.76 (0.56 to 1.04) |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | Stroke | Fosinopril Placebo | 18/196 11/201 | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | New focal neurological deficits of vascular origin with s/sx>24h, or death if occurred earlier | Telmisartan Placebo | 112/2954 136/2972 | AHR 0.83 (0.64 to 1.06) |

† = Enalapril vs placebo; ¶ = Amlodipine vs enalapril; ‡ = Amlodipine vs placebo; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; * = Clinical outcome data provided by FOSIDIAL corresponding author; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 6. KQ1 Composite – Cardiovascular Mortality, Nonfatal Myocardial Infarction, or Stroke – Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|------------------------------------|----------------------|-------------------------|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Ramipril Placebo | 651/4645 826/4652 | RR 0.78 (0.70 to 0.86) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) w/o HF | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | HTN and CAD | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Trandolapril Placebo | 396/4158 420/4132 | HR 0.93 (0.81 to 1.07) |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | Fosinopril Placebo | 48/196 41/201 | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Telmisartan Placebo | 384/2954 440/2972 | AHR 0.86 (0.74 to 1.00) |

† = Enalapril vs placebo; ‡ = Amlodipine vs placebo; ¶ = Amlodipine vs enalapril

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

* = Clinical outcome data provided by FOSIDIAL corresponding author

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 7. KQ1 Atrial Fibrillation - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|-------------------------|------------------------------------|----------------------|-------------------------|
| HOPE, 2000 ⁴⁰ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Atrial Fibrillation | Ramipril Placebo | 86/4291 91/4044 | OR 0.92 (0.68 to 1.24) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006* ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | New atrial fibrillation | Telmisartan Placebo | 182/2954 180/2972 | AHR 1.02 (0.83 to 1.26) |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 8. KQ1 Hospitalizations - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--|------------------------------------|------------------------|------------------------|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | NR | Ramipril Placebo | NR | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Admitted to the hospital at least once | Ramipril Placebo | 279/308 289/309 | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006* ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Number of patients hospitalized | Telmisartan Placebo | 1477/2954 1526/2972 | RR 0.97 (0.93 to 1.02) |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 9. KQ1 Hospitalization For Angina - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|---|------------------------------------|----------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Hospitalization for unstable angina | Ramipril Placebo | 554/4645 565/4652 | RR 0.98 (0.87 to 1.10) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Unstable angina requiring hospitalization | Ramipril Placebo | 45/308 42/309 | RR 1.08 (0.71 to 1.65) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Hospitalization for angina | Enalapril Placebo | 40/229 29/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Hospitalization for worsening angina | Candesartan Control | 9/194 14/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Hospitalization for angina | Enalapril Amlodipine Placebo | 86/673 51/663 84/655 | HR 0.98 (0.72 to 1.32) [†] HR 0.59 (0.42 to 0.84) [¶] HR 0.58 (0.41 to 0.82) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | Angina pectoris requiring hospitalization | ACEI [∞] Nifedipine | 56/822 50/828 | RR 0.80 (0.55 to 1.18) [∂] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006 ^{*48} | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Angina with hospitalization and ECG changes | Telmisartan Placebo | 253/2954 287/2972 | HR 0.88 (0.74 to 1.04) |

† = Enalapril vs placebo; ‡ = Amlodipine vs placebo; ¶ = Amlodipine vs enalapril; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 10. KQ1 Hospitalization For Heart Failure - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|---|------------------------------------|-------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Hospitalization for heart failure | Ramipril Placebo | 141/4645 160/4652 | RR 0.88 (0.70 to 1.10) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | CHF requiring hospitalization | Ramipril Placebo | 7/308 9/309 | RR 0.78 (0.29 to 2.09) |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | HF requiring hospital admission | Perindopril Placebo | 63/6110 103/6108 | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Hospitalization for HF | Candesartan Control | 0/194 2/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Hospitalization for CHF | Enalapril Amlodipine Placebo | 4/673 3/663 5/655 | HR 0.78 (0.21 to 2.90) [†] HR 0.78 (0.17 to 3.47) [¶] HR 0.59 (0.14 to 2.47) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | HF requiring hospitalization | ACEI [∞] Nifedipine | 9/822 12/828 | RR 1.25 (0.52 to 2.98) [∂] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | CHF as primary cause of hospitalization | Trandolapril Placebo | 105/4158 134/4132 | HR 0.77 (0.60 to 1.00) |
| FOSIDIAL, 2006* ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Hospitalization for HF or attendance in an acute care setting | Telmisartan Placebo | 134/2954 129/2972 | HR 1.05 (0.82 to 1.34) |

† = Enalapril vs placebo; ‡ = Amlodipine vs placebo; ¶ = Amlodipine vs enalapril; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 11. KQ1 Revascularization - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--|------------------------------------|-----------------------------|---|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | All CV revasc (CABG, PCI, carotid endarterectomy, peripheral vascular surgery) | Ramipril Placebo | 742/4645 852/4652 | RR 0.85 (0.77 to 0.94) |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | Any revascularization | Enalapril Placebo | 16/229 25/231 | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | Revasc (CABG or PTCA) | Perindopril Placebo | 577/6110 601/6108 | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | Revascularization | Candesartan Control | 8/194 15/203 | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Coronary revascularization | Enalapril Amlodipine Placebo | 95/673 78/663 103/655 | HR 0.86 (0.65 to 1.14) [†] HR 0.84 (0.62 to 1.13) [¶] HR 0.73 (0.54 to 0.98) [‡] |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | Performance of coronary interventions (PTCA, CABG or stenting) | ACEI [∞] Nifedipine | 75/822 81/828 | RR 1.04 (0.76 to 1.43) [⊙] |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | CABG | Trandolapril Placebo | 271/4158 294/4132 | HR 0.91 (0.77 to 1.07) |
| | | | PCI | Trandolapril Placebo | 515/4158 497/4132 | HR 1.03 (0.97 to 1.16) |
| FOSIDIAL, 2006* ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |

Appendix C: Additional Evidence Tables and Analyses


Appendix Table 11 Continued. KQ1 Revascularization - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------------|--------------|---|------------------------------|------------------------|----------------------|------------------------|
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Revascularization procedures | Telmisartan Placebo | 349/2954 390/2972 | HR 0.90 (0.77 to 1.03) |

† = Enalapril vs placebo; ‡ = Amlodipine vs placebo; ¶ = Amlodipine vs enalapril; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril; * = Clinical outcome data provided by FOSIDIAL corresponding author; ∂ = Nifedipine vs ACEI

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 1. KQ1 Total Mortality ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=2.064483 (df=5) p=0.8402

I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 2. KQ1 Total Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled & open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=9.913118 (df=7) p=0.1936
 I^2 statistic=29.4%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 3. KQ1 Total Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled & open-label trials in patients with stable ischemic heart disease risk equivalents




Test for heterogeneity: Cochran Q=4.461381 (df=1) p=0.0347

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 4. KQ1 Cardiovascular Mortality ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=7.343875 (df=4) p=0.1188

I^2 statistic=45.5%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 5. KQ1 Cardiovascular Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=14.733985 (df=6) p=0.0224

I^2 statistic=59.3%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 6. KQ1 Nonfatal Myocardial Infarction ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=7.189476$ (df=5) $p=0.2069$

I^2 statistic=30.5%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 7. KQ1 Nonfatal Myocardial Infarction Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=7.76543 (df=6) p=0.2558

I^2 statistic=22.7%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 8. KQ1 Stroke ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=8.03054 (df=5) p=0.1546
 I^2 statistic=37.7%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 9. KQ1 Stroke Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=8.291835 (df=6) p=0.011848
 I^2 statistic=27.6%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 10. KQ1 composite of Cardiovascular Mortality, Nonfatal Myocardial Infarction and Stroke ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=4.365658$ ($df=1$) $p=0.0367$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 11. KQ1 Hospitalization For Angina ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=2.371505$ (df=3) $p=0.499$

I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 12. KQ1 Hospitalization For Angina Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=4.876247$ (df=5) $p=0.4312$

I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 13. KQ1 Hospitalization For Heart Failure ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=3.530577$ (df=4) $p=0.4732$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 14. KQ1 Hospitalization For Heart Failure Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=8.660173$ (df=6) $p=0.1936$
 I^2 statistic=30.7%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 15. KQ1 Revascularization ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=2.989717$ (df=3) $p=0.3932$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 16. KQ1 Revascularization Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease undergoing coronary artery bypass grafting surgery only




Test for heterogeneity: Cochran $Q=0.037509$ ($df=1$) $p=0.8464$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 17. KQ1 Revascularization Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease undergoing percutaneous coronary intervention only




Test for heterogeneity: Cochran $Q=1.864482$ ($df=1$) $p=0.1721$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 18. KQ1 Revascularization Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease



Test for heterogeneity: Cochran $Q=4.252035$ ($df=5$) $p=0.5137$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 12. KQ3 Total Mortality - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|----------------------------------|--------------|--|-----------------------------|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | Death | Cilazapril Placebo | 7/1075 1/361 | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%; 5-7 days prior to randomization) or PTCA (18%; 1-2 days prior to randomization) for angina | Mortality due to all causes | Ramipril Placebo | 2/80 8/79 | 1-RR 76% (-1 to 92) |
| Kondo et al, 2001 ^{55†} | RCT | Received elective balloon angioplasty followed by coronary stenting | Total mortality | Quinapril Control | 0/49 0/50 | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | Deaths | Quinapril Placebo | 0/46 0/45 | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | All cause mortality | Quinapril Placebo | 27/878 27/872 | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | Deaths | Candesartan Placebo | 0/63 0/57 | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Death due to any cause | Quinapril Placebo | 28/1280 28/1273 | AHR 1.00 (0.59 to 1.69) |

† Outcomes provided by personal communication with corresponding author

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 13. KQ3 Cardiovascular Mortality - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|----------------------------------|--------------|--|---|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%; 5-7 days prior to randomization) or PTCA (18%; 1-2 days prior to randomization) for angina | CV death, including cardiac death and fatal stroke | Ramipril Placebo | 1/80 8/79 | 1-RR 88% (24 to 94) |
| Kondo et al, 2001 ^{55†} | RCT | Received elective balloon angioplasty followed by coronary stenting | Cardiovascular death | Quinapril Control | 0/49 0/50 | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | Deaths‡ | Quinapril Placebo | 0/46 0/45 | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | CV death, including cardiac death and vascular/stroke death | Quinapril Placebo | 13/878 14/872 | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | Deaths‡ | Candesartan Placebo | 0/63 0/57 | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Cardiovascular death | Quinapril Placebo | 18/1280 15/1273 | AHR 1.20 (0.60 to 2.38) |

† Outcomes provided by personal communication with corresponding author; ‡ No deaths occurred during the study.

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 14. KQ3 Nonfatal Myocardial Infarction - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|---|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | Nonfatal MI | Cilazapril Placebo | 27/1075 8/361 | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%; 5-7 days prior to randomization) or PTCA (18%; 1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | Nonfatal MI | Quinapril Placebo | 1/46 0/45 | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | Nonfatal MI defined as changes in 1 or more of three parameters: symptomatology, enzyme elevation and ECG changes | Quinapril Placebo | 36/878 40/872 | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | Nonfatal MI [†] | Candesartan Placebo | 1/63 2/57 | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Nonfatal MI | Quinapril Placebo | 16/1280 21/1273 | AHR 0.76 (0.40 to 1.46) |

[†] AACHEN reported no deaths in the trial, with one MI in the Candesartan group and two in the placebo group therefore events were entered as nonfatal.

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 15. KQ3 Stroke - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|--------------------|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Fatal stroke | Ramipril Placebo | 0/80 1/79 | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Stroke | Quinapril Placebo | 15/1280 14/1273 | AHR 1.07 (0.52 to 2.21) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 16. KQ3 Composite: Cardiovascular Mortality, Nonfatal Myocardial Infarction, or Stroke - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Ramipril Placebo | NR | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Quinapril Placebo | 45/1280 45/1273 | AHR 1.00 (0.66 to 1.51) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 17. KQ3 Atrial Fibrillation - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|----------------------------------|--------------|--|---|------------------------|----------------------|-----------------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo et al, 2001 ^{55†} | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | New-onset atrial fibrillation (after randomization) | Quinapril Placebo | 114/1280 101/1273 | % risk difference 1 (-1.2 to 3.1) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 18. KQ3 Hospitalization For Angina - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|---|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Patients hospitalized with chest pain on suspicion of unstable angina | Ramipril Placebo | 12/80 9/79 | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | Patients hospitalized with unstable angina | Quinapril Placebo | 45/878 52/872 | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Hospitalization for unstable angina | Quinapril Placebo | 45/1280 38/1273 | AHR 1.19 (0.77 to 1.83) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 19. KQ3 Hospitalization For Heart Failure - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**


| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|-----------------------------------|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Hospitalization for heart failure | Ramipril Placebo | 2/80 5/79 | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Hospitalization for heart failure | Quinapril Placebo | 15/1280 14/1273 | AHR 1.09 (0.53 to 2.26) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 20. KQ3 Revascularization - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|---------------------------------|--------------|--|-----------------------------------|------------------------|--------------------|-------------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | CABG or repeat angioplasty | Cilazapril Placebo | 207/1075 54/361 | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo et al, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | Angioplasty or stent implantation | Quinapril Placebo | 10/46 7/45 | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | Coronary angioplasty | Quinapril Placebo | 223/878 233/872 | NR |
| | | | CABG | Quinapril Placebo | 116/878 104/872 | |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | Target lesion revascularization | Candesartan Placebo | 5/63 4/57 | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Coronary revascularization | Quinapril Placebo | 52/1280 41/1273 | AHR 1.28 (0.85 to 1.93) |

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 19. KQ3 Total Mortality ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran Q=3.810035 (df=4) p=0.4323

I² statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 20. KQ3 Total Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=3.811901$ (df=6) $p=0.7021$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 21. KQ3 Total Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=3.810035$ ($df=4$) $p=0.4323$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 22. KQ3 Total Mortality Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, percutaneous procedure only




Test for heterogeneity: Cochran $Q=0.623781$ ($df=3$) $p=0.891$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 23. KQ3 Cardiovascular Mortality ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=4.351836$ (df=3) $p=0.2259$

I^2 statistic=31.1%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 24. KQ3 Cardiovascular Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=4.350679$ ($df=5$) $p=0.5001$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 25. KQ3 Cardiovascular Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=4.351836$ ($df=3$) $p=0.2259$
 I^2 statistic=31.1%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 26. KQ3 Cardiovascular Mortality Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, percutaneous procedure only




Test for heterogeneity: Cochran $Q=0.000956$ ($df=2$) $p=0.9995$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 27. KQ3 Nonfatal Myocardial Infarction ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.11656$ ($df=3$) $p=0.767$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 28. KQ3 Nonfatal Myocardial Infarction Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.46284$ (df=4) $p=0.8332$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 29. KQ3 Nonfatal Myocardial Infarction Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.141656$ (df=3) $p=0.767$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 30. KQ3 Nonfatal Myocardial Infarction Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, percutaneous procedure only




Test for heterogeneity: Cochran $Q=1.130237$ (df=3) $p=0.7698$

I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 31. KQ3 Stroke ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=0.497689$ ($df=1$) $p=0.4805$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 32. KQ3 Stroke Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=0.497689$ ($df=1$) $p=0.4805$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 33. KQ3 Stroke Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=0.497689$ ($df=1$) $p=0.4805$

I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 34. KQ3 Hospitalization For Angina ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.573147$ (df=2) $p=0.4554$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 35. KQ3 Hospitalization For Angina Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.573147$ (df=2) $p=0.4554$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 36. KQ3 Hospitalization For Angina Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=1.573147$ (df=2) $p=0.4554$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 37. KQ3 Revascularizations ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran Q=0.043777 (df=2) p=0.9783

I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 38. KQ3 Revascularizations Sensitivity Analysis - Meta-analysis of randomized placebo-controlled or open-label trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=0.082314$ ($df=3$) $p=0.9939$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 39. KQ3 Revascularizations Sensitivity Analysis - Meta-analysis of randomized placebo-controlled trials utilizing intention-to-treat methodologies in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, a coronary revascularization procedure




Test for heterogeneity: Cochran $Q=0.043777$ (df=2) $p=0.9783$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 40. KQ3 Revascularizations Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, percutaneous procedure only




Test for heterogeneity: Cochran $Q=4.040768$ ($df=3$) $p=0.2571$

I^2 statistic=25.8%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 41. KQ3 Revascularizations Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease who have recently undergone, or are set to undergo, coronary artery bypass grafting surgery only



Test for heterogeneity: Cochran $Q=0.291311$ ($df=1$) $p=0.5894$
 I^2 statistic=N/A

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 21. KQ4 Run-in Phase Data - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Run-in | Description | Exclusions |
|-------------------------------------|--------|---|--|
| HOPE, 2000 ³⁸ | Yes | Ramipril 2.5mg/d X 7-10d, then placebo qd X 10-14d | 1,035/10,576 (9.8%) excluded: <ul style="list-style-type: none"> - Non-compliance (n=NR) - ADE (n=NR) - Abnormal Scr or potassium (n=NR) - Withdrawal of consent (n=NR) |
| PART-2, 2000 ⁴¹ | Yes | Ramipril 5mg/d X 7d, then 10mg/d X 7d | 127/744 (17%) excluded: <ul style="list-style-type: none"> - Ineligibility (n=52, 41%) - Suspected ADE (n=52, 41%) - Patient preference (n=23, 18%) |
| SCAT, 2000 ⁴² | Yes | Dietary and SB placebo X 1 month | ~33% excluded [†] |
| EUROPA, 2003 ⁴³ | Yes | Perindopril 4mg/d X14d, then 8mg/d X 14d [‡] | 1,437/13,655 (10.5%) excluded: <ul style="list-style-type: none"> - Hypotension (n=290, 20.2%) - Raised Scr or potassium (n=149, 10.4%) - Other intolerance (n=332, 23.1%) - Major clinical event (n=75, 5.2%) - Poor adherence (n=80, 5.6%) - Exclusion criteria (n=44, 3.1%) - Withdrawn consent (n=9, 0.6%) - Unspecified stop reason (n=446, 31%) - Never randomized (n=12, 0.8%) |
| Kondo et al, 2003 ⁴⁴ | No | N/A | N/A |
| CAMELOT, 2004 ⁴⁵ | Yes | Placebo tablet + placebo capsule qd X 14d | NR |
| JMIC-B, 2004 ⁴⁶ | No | N/A | N/A |
| PEACE, 2004 ⁴⁷ | Yes | Trandolapril 2mg/d X 14d | NR |
| FOSIDIAL, 2006 ⁴⁸ | Yes | Single-blind placebo X 14d, then fosinopril 5mg X1 dose | NR |
| Takahashi et al, 2006 ⁴⁹ | No | N/A | N/A |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | No | N/A | N/A |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 21 Continued. KQ4 Run-in Phase Data - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Run-in | Description | Exclusions |
|----------------------------------|--------|---|---|
| TRANSCEND, 2008 ⁵¹ | Yes | Placebo qd X 7d, then Telmisartan 80mg/d X 14d | 740/6666 (11.1%) excluded: <ul style="list-style-type: none"> - Poor compliance (n=311, 42.0%) - Consent withdrawn (n=135, 18.2%) - Raised Scr or potassium (n=37, 5.0%) - Symptomatic hypotension (n=53, 7.2%) - Deaths (n=3, 0.4%) - Other reasons (n=201, 27.2%) |

† = Specific numbers of patients who entered run-in, and number who were excluded following run-in were not provided

‡ = Patients >70 years old received Perindopril 2mg/d X 7d, then 4mg/d X 7d, then 8mg/d X 14d during run-in period

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 22. KQ4 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Report Withdrawals | Group | n | Reasons |
|----------------------------|--------------------|-----------|--------------|--|
| HOPE, 2000 ³⁸ | Yes | Ramipril | 1511 (32.5%) | Cough (n=340, 7.3%) Hypotension/Dizziness (n=88, 1.9%) Angioedema (n=17, 0.4%) Uncontrolled HTN (n=109, 2.3%) Clinical Events (n=309, 6.7%) Other (n=1101, 23.7%) |
| | | Placebo | 1430 (30.7%) | Cough (n=85, 1.8%) Hypotension/Dizziness (n=70, 1.5%) Angioedema (n=7, 0.2%) Uncontrolled HTN (n=183, 3.9%) Clinical Events (n=418, 9.0%) Other (n=1074, 23.1%) |
| PART-2, 2000 ⁴¹ | Yes | Ramipril | 53 (17.2%) | Suspected ADE (n=31, 10%) Patient preference (n=22, 7%) |
| | | Placebo | 25 (8.1%) | Suspected ADE (n=3, 1%) Patient preference (n=22, 7%) |
| SCAT, 2000 ⁴² | No | Enalapril | N/A | N/A |
| | | Placebo | | |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 22 Continued. KQ4 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Report Withdrawals | Group | n | Reasons |
|---------------------------------|--------------------|-------------|--------------|--|
| EUROPA, 2003 ⁴³ | Yes | Perindopril | 1391 (22.8%) | Cough (n=162, 2.7%) Hypotension (n=60, 1.0%) Kidney failure (n=20, 0.3%) Intolerance (n=144, 2.4%) Study endpoint (n=376, 6.2%) Hypertension (n=22, 0.4%) Refusal to continue (n=261, 4.3%) Other (n=347, 5.7%) |
| | | Placebo | 1266 (20.7%) | Cough (n=32, 0.5%) Hypotension (n=17, 0.3%) Kidney failure (n=16, 0.3%) Intolerance (n=80, 1.3%) Study endpoint (n=460, 7.5%) Hypertension (n=46, 0.8%) Refusal to continue (n=257, 4.2%) Other (n=359, 5.9%) |
| Kondo et al, 2003 ⁴⁴ | Yes | Candesartan | 9 (4.4%) | Dizziness/Lightheadedness (n=9, 4.4%) |
| | | Control | 2 (1.0%) | Relocation (n=2, 1.0%) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 22 Continued. KQ4 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Report Withdrawals | Group | n | Reasons |
|-----------------------------|--------------------|------------|-------------|---|
| CAMELOT, 2004 ⁴⁵ | Yes | Enalapril | 236 (35.0%) | ADE (n=102, 15.1%) Withdrew consent (n=33, 4.9%) Death (n=4, 0.6%) Protocol violation (n=6, 0.9%) Laboratory abnormality (n=3, 0.4%) Lost to follow-up (n=22, 3.3%) Insufficient response (n=5, 0.7%) Other (n=61, 9.0%) |
| | | Amlodipine | 194 (29.2%) | ADE (n=87, 13.1%) Withdrew consent (n=38, 5.7%) Death (n=2, 0.3%) Laboratory abnormality (n=2, 0.3%) Lost to follow-up (n=18, 2.7%) Insufficient response (n=2, 0.3%) Other (n=45, 6.8%) |
| | | Placebo | 204 (31.1%) | ADE (n=71, 10.8%) Withdrew consent (n=50, 7.6%) Death (n=5, 0.8%) Protocol violation (n=8, 1.2%) Laboratory abnormality (n=3, 0.5%) Lost to follow-up (n=16, 2.4%) Insufficient response (n=3, 0.5%) Other (n=48, 7.3%) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 22 Continued. KQ4 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Report Withdrawals | Group | n | Reasons |
|-------------------------------------|--------------------|-------------------|-------------|--|
| JMIC-B, 2004 ⁴⁶ | Yes | ACEI [∞] | 143 (17.4%) | ADE (n=72, 8.8%) No effect (n=20, 2.4%) Withdrawal of consent (n=10, 1.2%) Protocol deviation (n=5, 0.6%) Alleviating symptoms (n=11, 1.3%) Others (n=25, 3.0%) |
| | | Nifedipine | 107 (12.9%) | ADE (n=41, 5.0%) No effect (n=11, 1.3%) Withdrawal of consent (n=9, 1.1%) Protocol deviation (n=9, 1.1%) Alleviating symptoms (n=17, 2.1%) Others (n=20, 2.4%) |
| PEACE, 2004 ^{†47} | No | Trandolapril | N/A | N/A |
| | | Placebo | | |
| FOSIDIAL, 2006 ⁴⁸ | Yes | Fosinopril | 7 (3.6%) | Renal transplantation (n=7, 3.6%) |
| | | Placebo | 10 (5.0%) | Renal transplantation (n=8, 4.0%) Protocol violations (n=2, 1.0%) |
| Takahashi et al, 2006 ⁴⁹ | Yes | Candesartan | 0 (0%) | N/A |
| | | Control | 0 (0%) | |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | Yes | Zofenopril | 46 (13.2%) | Major protocol violation (n=15, 4.3%) Lost to follow-up (n=3, 0.9%) Inability to perform treadmill test (n=31, 8.9%) |
| | | Placebo | | |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

† = Listed as side effects leading to discontinuation of the study medication

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 22 Continued. KQ4 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Report Withdrawals | Group | n | Reasons |
|-------------------------------|--------------------|-------------|--------------|--|
| TRANSCEND, 2008 ⁵¹ | Yes | Telmisartan | 1090 (36.9%) | Hypotensive symptoms (n=29, 1.0%) Syncope (n=1, 0.03%) Cough (n=15, 0.5%) Diarrhea (n=7, 0.2%) Angioedema (n=2, 0.07%) Renal abnormalities (n=24, 0.8%) |
| | | Placebo | 1143 (38.5%) | Hypotensive symptoms (n=16, 0.5%) Syncope (n=0, 0%) Cough (n=18, 0.6%) Diarrhea (n=2, 0.07%) Angioedema (n=3, 0.1%) Renal abnormalities (n=13, 0.4%) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 23. KQ4 Withdrawals Due To Adverse Events - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|---|------------------------------------|-----------------------------|-----------------------|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | NR | Ramipril Placebo | NR | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | Suspected adverse drug reactions leading to stopping randomized treatment | Ramipril Placebo | 31/308 3/309 | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Discontinuations due to adverse events | Enalapril Amlodipine Placebo | 102/673 87/663 71/655 | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | Withdrawals due to adverse events | ACEI [∞] Nifedipine | 72/822 41/828 | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Side effects leading to discontinuation of study medication | Trandolapril Placebo | 599/4158 269/4132 | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 23 Continued. KQ4 Withdrawals Due To Adverse Events - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease.**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------|--------------|---|--------------------|------------------------|-------------|-----------------------|
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 24. KQ4 Hypotension - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|----------------------------|-----------------------|
| HOPE, 2000 ^{38†} | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Hypotension | Ramipril Placebo | 2/4645 3/4652 | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Hypotension | Enalapril Amlodipine Placebo | 64/673 22/663 21/655 | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | Severe hypotension | Zofenopril Placebo | 2/172 2/177 | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

† = Data are reported as “serious adverse events” found within the New Drug Application from www.fda.gov.

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 25. KQ4 Syncope - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|----------------------|-----------------------|
| HOPE, 2000 ^{38†} | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Syncope | Ramipril Placebo | 3/4645 1/4652 | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Syncope | Trandolapril Placebo | 200/4158 161/4132 | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

† = Data are reported as “serious adverse events” found within the New Drug Application from www.fda.gov.

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 26. KQ4 Cough - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|----------------------------|-----------------------|
| HOPE, 2000 ^{38†} | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Cough | Ramipril Placebo | 16/4645 9/4652 | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | Cough | Enalapril Amlodipine Placebo | 84/673 34/663 38/655 | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Cough | Trandolapril Placebo | 1626/4158 1136/4132 | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

† = Data are reported as “serious adverse events” found within the New Drug Application from www.fda.gov.

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 27. KQ4 Angioedema - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|------------------|-----------------------|
| HOPE, 2000 ^{38†} | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Angioedema | Ramipril Placebo | 5/4645 1/4652 | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | Angioedema | Trandolapril Placebo | 8/4158 5/4132 | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

† = Data are reported as “serious adverse events” found within the New Drug Application from www.fda.gov.

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 28. KQ4 Hyperkalemia - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|------------------------------------|------------------------------------|----------------------|-----------------------|
| HOPE, 2005 ^{38†} | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | Serum potassium level > 5.0 mmol/L | Ramipril Placebo | 395/4539 297/4572 | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | Serum potassium level > 5.5 mmol/L | Telmisartan Placebo | 111/2954 49/2972 | NR |

† = Data taken from Mann JFE, et al. Serum potassium, cardiovascular risk, and effects of an ACE inhibitor: results of the HOPE Study. Clin Nephrol 2005;63:181-7; ∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 29. KQ4 Blood Dyscrasias - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|-------------|-----------------------|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | NR | Ramipril Placebo | NR | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril


Appendix C: Additional Evidence Tables and Analyses**Appendix Table 30. KQ4 Rash - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|-------------------------------------|--------------|--|--------------------|------------------------------------|-------------|-----------------------|
| HOPE, 2000 ³⁸ | RCT | CAD, Stroke, PVD or DM + 1 Risk Factor | NR | Ramipril Placebo | NR | NR |
| PART-2, 2000 ⁴¹ | RCT | MI, angina with confirmed CAD, TIA or IC | NR | Ramipril Placebo | NR | NR |
| SCAT, 2000 ⁴² | RCT | Coronary atherosclerosis in >3 major arteries, elevated cholesterol | NR | Enalapril Placebo | NR | NR |
| EUROPA, 2003 ⁴³ | RCT | CAD (previous MI, revasc. or >70% coronary artery narrowing) without HF | NR | Perindopril Placebo | NR | NR |
| Kondo et al, 2003 ⁴⁴ | RCT | H/o coronary intervention with no significant stenosis on 6 mo f/o angiography | NR | Candesartan Control | NR | NR |
| CAMELOT, 2004 ⁴⁵ | RCT | PCI or chest pain requiring coronary angiography | NR | Enalapril Amlodipine Placebo | NR | NR |
| JMIC-B, 2004 ⁴⁶ | RCT | Hypertension and CAD | NR | ACEI [∞] Nifedipine | NR | NR |
| PEACE, 2004 ⁴⁷ | RCT | Documented CAD | NR | Trandolapril Placebo | NR | NR |
| FOSIDIAL, 2006 ⁴⁸ | RCT | Hemodialysis and LVH | NR | Fosinopril Placebo | NR | NR |
| Takahashi et al, 2006 ⁴⁹ | RCT | Chronic maintenance hemodialysis | NR | Candesartan Control | NR | NR |
| SMILE-ISCHEMIA, 2007 ⁵⁰ | RCT | MI within 6 weeks | NR | Zofenopril Placebo | NR | NR |
| TRANSCEND, 2008 ⁵¹ | RCT | CAD, Cerebrovascular disease, PVD, or DM + end-organ damage | NR | Telmisartan Placebo | NR | NR |

∞ = Patients in the ACEI group were given enalapril, imidapril, or lisinopril

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 42. KQ4 Withdrawal Due To Adverse Events Subgroup ACEI Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran Q=15.650446 (df=2) p=0.0004
 I^2 statistic=87.2%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 43. KQ4 Hypotension ACEI Subgroup Analysis - Meta-analysis of randomized placebo-controlled trials in patients with stable ischemic heart disease



Test for heterogeneity: Cochran $Q=3.368646$ ($df=2$) $p=0.1856$

I^2 statistic=40.6%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 31. KQ6 Run-in Phase Data - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease

| Study, year | Run-in | Description | Exclusions |
|---------------------------------|--------|-------------|------------|
| MARCATOR, 1995 ⁵³ | No | N/A | N/A |
| APRES, 2000 ⁵⁴ | No | N/A | N/A |
| Kondo et al, 2001 ⁵⁵ | No | N/A | N/A |
| PARIS, 2001 ⁵⁶ | No | N/A | N/A |
| QUIET, 2001 ⁵⁷ | No | N/A | N/A |
| AACHEN, 2006 ⁵⁸ | No | N/A | N/A |
| IMAGINE, 2008 ⁵⁹ | No | N/A | N/A |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 32. KQ6 Study Withdrawals - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Report Withdrawals | Group | n | Reasons |
|------------------------------|--------------------|--------------------------|--------------------------------|--|
| MARCATOR, 1995 ⁵³ | Yes | Cilazapril Placebo | 350 (24.4%) [†] | No follow-up angiogram (n=159, 11.1%) Protocol violation (n=22, 1.5%) Severe hypotension (n=33, 2.3%) Severe cough (n=21, 1.5%) Angina pectoris (range 10-14% per group) |
| APRES, 2000 ⁵⁴ | Yes | Ramipril Placebo | 13 (16.3%) 13 (16.5%) | Open-label ACEI treatment (n=5, 6.3%) Loss of consent/follow-up (n=5, 6.3%) Side effects (n=2, 2.5%) Endocarditis requiring surgery (n=1, 1.3%) Open-label ACEI treatment (n=7, 8.9%) Loss of consent/follow-up (n=4, 5.1%) Side effects (n=2, 2.5%) |
| Kondo, 2001 ⁵⁵ | Yes | Quinapril Control | 1 (2%) 0 (0%) | Severe cough (n=1, 2%) N/A |
| PARIS, 2001 ⁵⁶ | Yes | Quinapril Placebo | 0 (0%) 0 (0%) | N/A 0 (0%) |
| QUIET, 2001 ⁵⁷ | Yes | Quinapril Placebo | 246 (28.0%) 218 (25.0%) | NR NR |
| AACHEN, 2006 ⁵⁸ | No | Candesartan Placebo | N/A | N/A |
| IMAGINE, 2008 ⁵⁹ | Yes | Quinapril Placebo | 444 (34.7%) 321 (25.2%) | Adverse event (n=228, 17.8%) Worsening diabetes (n=8, 0.6%) Patient decision (n=103, 8.0%) Physician decision (n=73, 5.7%) Other (n=32, 2.5%) Adverse event (n=103, 8.1%) Worsening diabetes (n=3, 0.2%) Patient decision (n=89, 7.0%) Physician decision (n=97, 7.6%) Other (n=23, 1.8%) |

† = Reasons for all 350 patient withdrawals was not given

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 33. KQ6 Withdrawals Due To Adverse Events - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--|------------------------|----------------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Withdrawal due to side effects | Ramipril Placebo | 2/80 2/79 | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | Dropped out of the study due to adverse events | Quinapril Control | 1/49 0/50 | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | N/A [†] | Quinapril Placebo | 0/46 0/45 | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Discontinuations due to adverse events | Quinapril Placebo | 228/1280 103/1273 | NR |

[†] All patients were followed-up

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 34. KQ6 Hypotension - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|---------------------|---|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Hypotension | Quinapril Placebo | 154/1280 70/1273 | Absolute difference 6.5% (4.5% to 8.5%) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 35. KQ6 Syncope - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|-------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | NR | Quinapril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 36. KQ6 Cough - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|----------------------|---|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | Severe cough | Quinapril Control | 1/49 0/50 | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | Cough | Quinapril Placebo | 33/878 2/872 | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | Cough | Quinapril Placebo | 269/1280 141/1273 | Absolute difference 10% (7.2% to 12.7%) |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 37. KQ6 Angioedema - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|-------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | NR | Quinapril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 38. KQ6 Hyperkalemia - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------------|------------------------|--------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | Electrolytic derangement | Ramipril Placebo | 0/80 0/79 | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | NR | Quinapril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 39. KQ6 Rash - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**


| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|-------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | NR | Quinapril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses**Appendix Table 40. KQ6 Blood Dyscrasias - Comparative effectiveness of medical therapies with or without ACEI or ARBs for stable ischemic heart disease**

| Study, year | Study Design | Population | Outcome/Definition | Group | Events, n/N | Events, “X”R (95% CI) |
|------------------------------|--------------|--|--------------------|------------------------|-------------|-----------------------|
| MARCATOR, 1995 ⁵³ | RCT | Undergoing elective coronary angioplasty | NR | Cilazapril Placebo | NR | NR |
| APRES, 2000 ⁵⁴ | RCT | Underwent elective CABG (82%)(5-7 days prior to randomization) or PTCA (18%)(1-2 days prior to randomization) for angina | NR | Ramipril Placebo | NR | NR |
| Kondo, 2001 ⁵⁵ | RCT | Received elective balloon angioplasty followed by coronary stenting | NR | Quinapril Control | NR | NR |
| PARIS, 2001 ⁵⁶ | RCT | Underwent successful elective PCI with stent implantation | NR | Quinapril Placebo | NR | NR |
| QUIET, 2001 ⁵⁷ | RCT | Underwent successful elective coronary angioplasty of atherectomy within 12-72 hours | NR | Quinapril Placebo | NR | NR |
| AACHEN, 2006 ⁵⁸ | RCT | Undergoing elective coronary stent implantation (treatment started 7-14 days prior to intervention) | NR | Candesartan Placebo | NR | NR |
| IMAGINE, 2008 ⁵⁹ | RCT | Underwent CABG (7-10 days prior) | NR | Quinapril Placebo | NR | NR |

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 44. KQ6. Withdrawals Due To Adverse Events Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease




Test for heterogeneity: Cochran $Q=0.856866$ ($df=3$) $p=0.8358$
 I^2 statistic=0%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Figure 45. KQ6. Cough Sensitivity Analysis - Meta-analysis of randomized placebo-controlled + open-label trials in patients with stable ischemic heart disease



Test for heterogeneity: Cochran $Q=9.185671$ ($df=2$) $p=0.0101$
 I^2 statistic=78.2%

Note: The squares represent individual point estimates. The size of the square represents the weight given to each study in the meta-analysis. Horizontal lines through each square represent 95% confidence intervals. The diamond represents the combined results. The solid vertical line extending from 1 is the null value.

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 41. KQ1 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | Evidence Grade | Importance |
|--|------------------|------------------------|--------------------------|-------------------------|------------------------|----------------------|---------------------|-------------------|------------------------|--|----------------|------------|
| | | | | | | | No of patients | | Effect | | | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Treatment | Control* | Relative Risk (95% CI) | Absolute Risk | | |
| Total mortality - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | |
| 7 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 1552/19077 (8.1%) | 1714/19059 (9%) | RR 0.91 (0.84 to 0.98) | 8 fewer per 1000 (from 2 fewer to 14 fewer) | HIGH | CRITICAL |
| | | | | | | | | 0.9% | | 0 fewer per 1,000 | | |
| | | | | | | | | 12% | | 10 fewer per 1,000 | | |
| Total mortality - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 23/1495 (1.5%) | 19/1491 (1.3%) | RR 1.21 (0.66 to 2.21) | 3 more per 1000 (from 4 fewer to 16 more) | MODERATE | CRITICAL |
| | | | | | | | | 1.1% | | 2 more per 1,000 | | |
| | | | | | | | | 1.4% | | 2 more per 1,000 | | |
| Total mortality - IHD risk equivalents (follow-up 1.6-4.8 years) | | | | | | | | | | | | |
| 1 | randomized trial | single trial | no serious inconsistency | no serious indirectness | very serious | none | 53/196 (27.0%) | 50/201 (24.9%) | RR 1.08 (0.78 to 1.52) | 158 fewer per 1000 (from 234 fewer to 1000 more) | LOW | CRITICAL |
| Cardiovascular mortality - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | |
| 6 | randomized trial | no serious limitations | serious | no serious indirectness | serious | none | 883/18848 (4.7%) | 1021/18828 (5.4%) | RR 0.87 (0.75 to 1.02) | 7 fewer per 1000 (from 14 fewer to 1 more) | LOW | CRITICAL |
| | | | | | | | | 0.3% | | 0 fewer per 1,000 | | |
| | | | | | | | | 8.1% | | 10 fewer per 1,000 | | |
| Cardiovascular mortality - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 11/1495 (0.7%) | 11/1491 (0.7%) | RR 1.00 (0.43 to 2.29) | 0 fewer per 1000 (from 4 fewer to 9 more) | MODERATE | CRITICAL |
| | | | | | | | | 0.73% | | 0 fewer per 1,000 | | |
| | | | | | | | | 0.75% | | 0 fewer per 1,000 | | |
| | | | | | | | | | | | | |

Appendix C: Additional Evidence Tables and Analyses

| Cardiovascular mortality - IHD risk equivalents (follow-up 4.8 years) | | | | | | | | | | | | |
|---|------------------|------------------------|--------------------------|-------------------------|------------------------|------|--------------------|--------------------|------------------------|--|----------|----------|
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 32/196 (16.3%) | 31/201 (15.4%) | RR 1.06 (0.67 to 1.67) | 9 more per 1000 (from 51 fewer to 103 more) | MODERATE | CRITICAL |
| Nonfatal myocardial infarction - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | |
| 6 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 813/16123 (5%) | 981/16087 (6.1%) | RR 0.83 (0.73 to 0.94) | 10 fewer per 1000 (from 4 fewer to 16 fewer) | HIGH | CRITICAL |
| | | | | | | | | 2.9% | | 4 fewer per 1,000 | | |
| | | | | | | | | 7.2% | | 12 fewer per 1,000 | | |
| Nonfatal myocardial infarction - vs. CCB (follow-up 2 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 11/673 (1.6%) | 14/663 (2.1%) | RR 0.77 (0.35 to 1.69) | 5 fewer per 1000 (from 14 fewer to 14 more) | MODERATE | CRITICAL |
| Nonfatal myocardial infarction - IHD risk equivalents (follow-up 4.8 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | very serious | none | 9/196 (4.6%) | 7/201 (3.5%) | RR 1.31 (0.50 to 3.47) | 11 more per 1000 (from 18 fewer to 86 more) | LOW | CRITICAL |
| Stroke - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | |
| 7 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 454/19077 (2.4%) | 581/19059 (3%) | RR 0.79 (0.67 to 0.93) | 6 fewer per 1000 (from 2 fewer to 10 fewer) | HIGH | CRITICAL |
| | | | | | | | | 1.3% | | 2 fewer per 1,000 | | |
| | | | | | | | | 4.9% | | 10 fewer per 1,000 | | |
| Stroke - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 24/1495 (1.6%) | 22/1491 (1.5%) | RR 1.09 (0.61 to 1.94) | 1 more per 1000 (from 6 fewer to 14 more) | MODERATE | CRITICAL |
| | | | | | | | | 0.9% | | 0 more per 1,000 | | |
| | | | | | | | | 1.9% | | 1 more per 1,000 | | |
| Stroke - IHD risk equivalents (follow-up 4.8 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | very serious | none | 18/196 (9.2%) | 11/201 (5.5%) | RR 1.60 (0.81 to 3.46) | 33 more per 1000 (from 10 fewer to 135 more) | LOW | CRITICAL |
| Cardiovascular mortality, myocardial infarction, stroke - IHD (follow-up 4.5-4.8 years) | | | | | | | | | | | | |
| 3 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 1431/11757 (12.2%) | 1686/11756 (14.3%) | RR 0.86 (0.77 to 0.95) | 20 fewer per 1000 (from 7 fewer to 33 fewer) | HIGH | CRITICAL |
| | | | | | | | | 10% | | 13 fewer per 1,000 | | |

Appendix C: Additional Evidence Tables and Analyses

| | | | | | | | | | | | | | |
|--|------------------|------------------------|--------------------------|-------------------------|------------------------|------|-------------------|-------------------|------------------------|---|--|-----------|-----------|
| | | | | | | | | 18% | | 25 fewer per 1,000 | | | |
| Cardiovascular mortality, myocardial infarction, stroke - IHD risk equivalents (follow-up 4.8 years) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | very serious | none | | 48/196 (24.5%) | 41/201 (20.4%) | RR 1.20 (0.83 to 1.73) | 41 more per 1000 (from 35 fewer to 149 more) | MODERATE | CRITICAL |
| Atrial fibrillation - IHD (follow-up 4.5-4.7 years) | | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 268/7245 (3.7%) | 271/7016 (3.9%) | RR 0.98 (0.83 to 1.15) | 1 fewer per 1000 (from 7 fewer to 6 more) | HIGH | IMPORTANT | |
| | | | | | | | | 2.3% | | 0 fewer per 1,000 | | | |
| | | | | | | | | 6.1% | | 1 fewer per 1,000 | | | |
| Angina symptoms: Treadmill exercise test (follow-up 6 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | | 151 | 152 | WMD 3.5 minutes | 3.5 (2.82 to 4.18) | MODERATE | IMPORTANT |
| Total Hospitalizations – IHD (follow-up 4.7 years) | | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 1756/3262 (53.8%) | 1815/3281 (55.3%) | RR 0.97 (0.94 to 1.00) | 17 fewer per 1000 (from 33 fewer to 0 more) | MODERATE | IMPORTANT | |
| | | | | | | | | 51% | | 15 fewer per 1,000 | | | |
| | | | | | | | | 94% | | 28 fewer per 1,000 | | | |
| Hospitalizations for angina - IHD (follow-up 2-4.7 years) | | | | | | | | | | | | | |
| 5 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 978/8809 (11.1%) | 1007/8819 (11.4%) | RR 0.97 (0.89 to 1.06) | 3 fewer per 1000 (from 13 fewer to 7 more) | HIGH | IMPORTANT | |
| | | | | | | | | 9.7% | | 2 fewer per 1,000 | | | |
| | | | | | | | | 13.6% | | 4 fewer per 1,000 | | | |
| Hospitalizations for angina - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 142/1498 (9.5%) | 101/1491 (6.8%) | RR 1.38 (0.95 to 2.02) | 26 more per 1000 (from 3 fewer to 69 more) | MODERATE | IMPORTANT | |
| | | | | | | | | 6% | | 22 more per 1,000 | | | |
| | | | | | | | | 7.7% | | 29 more per 1,000 | | | |
| Hospitalizations for heart failure - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | | |
| 6 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 454/18848 (2.4%) | 540/18828 (2.9%) | RR 0.83 (0.70 to 0.98) | 5 fewer per 1000 (from 1 fewer to 9 fewer) | HIGH | IMPORTANT | |
| | | | | | | | | 0.8% | | 1 fewer per 1,000 | | | |
| | | | | | | | | 4.3% | | 7 fewer per 1,000 | | | |
| Hospitalizations for heart failure - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | | |

Appendix C: Additional Evidence Tables and Analyses

| | | | | | | | | | | | | |
|--|------------------|------------------------|--------------------------|-------------------------|------------------------|------|--------------------|--------------------|------------------------|--|----------|-----------|
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 13/1495 (0.9%) | 15/1491 (1%) | RR 0.87 (0.41 to 1.83) | 1 fewer per 1000 (from 6 fewer to 8 more) | MODERATE | IMPORTANT |
| | | | | | | | | 0.5% | | 0 fewer per 1,000 | | |
| | | | | | | | | 1.4% | | 1 fewer per 1,000 | | |
| Need for revascularization - IHD (follow-up 2-4.7 years) | | | | | | | | | | | | |
| 5 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 1779/14611 (12.2%) | 1971/14618 (13.5%) | RR 0.90 (0.85 to 0.96) | 14 fewer per 1000 (from 5 fewer to 20 fewer) | HIGH | CRITICAL |
| | | | | | | | | 9.8% | | 9 fewer per 1,000 | | |
| | | | | | | | | 18.3% | | 18 fewer per 1,000 | | |
| Need for revascularization - vs. CCB (follow-up 2-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 170/1495 (11.4%) | 159/1491 (10.7%) | RR 1.06 (0.83 to 1.36) | 6 more per 1000 (from 18 fewer to 39 more) | MODERATE | CRITICAL |
| | | | | | | | | 9.8% | | 5 more per 1,000 | | |
| | | | | | | | | 11.8% | | 7 more per 1,000 | | |

*Risk in the control group is reported as observed pooled, low and high-risks derived from included trials

Abbreviations: CCB=calcium channel blocker; CI=confidence interval; CV=cardiovascular; IHD=ischemic heart disease; MI=myocardial infarction; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 42. KQ2 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | | Evidence Grade | Importance |
|---|------------------|------------------------|---------------|-------------------------|------------------------|----------------------|---------------------|-------------------|------------------------|---|----------|----------------|------------|
| | | | | | | | No of patients | | Effect | | | | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Treatment | Control | Relative Risk (95% CI) | Absolute Risk | | | |
| Total mortality (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 1065/8502 (12.5%) | 1014/8576 (11.8%) | RR 1.07 (0.98 to 1.16) | 8 more per 1000 (from 2 fewer to 19 more) | MODERATE | CRITICAL | |
| Cardiovascular mortality (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 620/8502 (7.3%) | 603/8576 (7%) | RR 1.04 (0.93 to 1.17) | 3 more per 1000 (from 5 fewer to 12 more) | MODERATE | CRITICAL | |
| Mmyocardial infarction (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 438/8502 (5.2%) | 413/8576 (4.8%) | RR 1.08 (0.94 to 1.23) | 4 more per 1000 (from 3 fewer to 11 more) | MODERATE | CRITICAL | |
| Stroke (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 373/8502 (4.4%) | 405/8576 (4.7%) | RR 0.93 (0.81 to 1.07) | 3 fewer per 1000 (from 9 fewer to 3 more) | MODERATE | CRITICAL | |
| Cardiovascular mortality, myocardial infarction, stroke (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | no serious imprecision | none | 1200/8502 (14.1%) | 1210/8576 (14.1%) | RR 1.00 (0.93 to 1.09) | 0 fewer per 1000 (from 10 fewer to 13 more) | HIGH | CRITICAL | |
| New onset atrial fibrillation (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 537/8502 (6.3%) | 570/8576 (6.6%) | RR 0.96 (0.85 to 1.07) | 3 fewer per 1000 (from 10 fewer to 5 more) | MODERATE | IMPORTANT | |
| Worsening/new angina (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 538/8502 (6.3%) | 567/8576 (6.6%) | RR 0.96 (0.85 to 1.08) | 3 fewer per 1000 (from 10 fewer to 5 more) | MODERATE | IMPORTANT | |
| Hospitalization for angina (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 952/8502 (11.2%) | 925/8576 (10.8%) | RR 1.04 (0.95 to 1.14) | 4 more per 1000 (from 5 fewer to 15 more) | MODERATE | IMPORTANT | |
| Hospitalization for heart failure (follow-up 56 months) | | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | serious | none | 332/8502 (3.9%) | 354/8576 (4.1%) | RR 0.95 (0.82 to 1.1) | 2 fewer per 1000 (from 7 fewer to 4 more) | MODERATE | CRITICAL | |
| | | | | | | | | | | | | | |

Appendix C: Additional Evidence Tables and Analyses

| Revascularization (follow-up 56 months) | | | | | | | | | | | | |
|---|------------------|------------------------|--------------|-------------------------|------------------------|------|-------------------|-------------------|------------------------|---|----------|----------|
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | no serious imprecision | none | 1303/8502 (15.3%) | 1269/8576 (14.8%) | RR 1.04 (0.97 to 1.13) | 6 more per 1000 (from 4 fewer to 19 more) | Moderate | CRITICAL |

Abbreviations: CI=confidence interval; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 43. KQ3 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | Evidence Grade | Importance |
|---|------------------|------------------------|--------------------------|-------------------------|-------------|----------------------|---------------------|-----------------|------------------------|---|----------------|------------|
| | | | | | | | No of patients | | Effect | | | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Treatment | Control* | Relative Risk (95% CI) | Absolute Risk | | |
| Total mortality (follow-up 0.5-3 years) | | | | | | | | | | | | |
| 6 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 64/3422 (1.9%) | 64/2687 (2.4%) | RR 0.94 (0.67 to 1.37) | 1 fewer per 1000 (from 8 fewer to 9 more) | MODERATE | CRITICAL |
| | | | | | | | | 0% | | 0 fewer per 1,000 | | |
| | | | | | | | | 10% | | 6 fewer per 1,000 | | |
| Cardiovascular mortality (follow-up 0.5-3 years) | | | | | | | | | | | | |
| 5 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 32/2347 (1.4%) | 37/2326 (1.6%) | RR 0.91 (0.53 to 1.57) | 1 fewer per 1000 (from 8 fewer to 9 more) | MODERATE | CRITICAL |
| | | | | | | | | 0% | | 0 fewer per 1,000 | | |
| | | | | | | | | 10% | | 8 fewer per 1,000 | | |
| Nonfatal myocardial infarction (follow-up 0.5-3 years) | | | | | | | | | | | | |
| 5 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 81/3342 (2.4%) | 71/2608 (2.7%) | RR 0.89 (0.65 to 1.24) | 3 fewer per 1000 (from 9 fewer to 6 more) | MODERATE | CRITICAL |
| | | | | | | | | 0% | | 0 fewer per 1,000 | | |
| | | | | | | | | 4.6% | | 5 fewer per 1,000 | | |
| Stroke (follow-up 2.8-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | serious | none | 15/1360 (1.1%) | 15/1352 (1.1%) | RR 1.01 (0.50 to 2.04) | 0 fewer per 1000 (from -1 fewer to 1 more) | MODERATE | CRITICAL |
| | | | | | | | | 1.1% | | 0 more per 1,000 | | |
| | | | | | | | | 1.3% | | 0 more per 1,000 | | |
| Cardiovascular mortality, myocardial infarction, stroke (follow-up 3 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | serious | none | 45/1280 (3.5%) | 45/1273 (3.5%) | RR 0.99 (0.66 to 1.49) | 0 fewer per 1000 (from 12 fewer to 17 more) | MODERATE | CRITICAL |
| Atrial fibrillation (follow-up 3 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | serious | none | 114/1280 (8.9%) | 101/1273 (7.9%) | RR 1.12 (0.87 to 1.45) | 9 more per 1000 (from 10 fewer to 36 more) | MODERATE | IMPORTANT |
| | | | | | | | | | | | | |

Appendix C: Additional Evidence Tables and Analyses

| Hospitalization for angina (follow-up 2.3-3 years) | | | | | | | | | | | | |
|---|------------------|------------------------|--------------------------|-------------------------|------------------------|------|------------------|-----------------|------------------------|--|------|-----------|
| 3 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 102/2238 (4.6%) | 99/2224 (4.5%) | RR 1.02 (0.78 to 1.34) | 1 more per 1000 (from 10 fewer to 15 more) | HIGH | IMPORTANT |
| | | | | | | | | 3.0% | | 0 more per 1,000 | | |
| | | | | | | | | 11.4% | | 2 more per 1,000 | | |
| Hospitalization for heart failure (follow-up 3 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | very serious | none | 15/1280 (1.2%) | 14/1273 (1.1%) | RR 1.07 (0.52 to 2.20) | 1 more per 1000 (from 5 fewer to 13 more) | LOW | CRITICAL |
| Revascularization (follow-up 0.5-3 years) | | | | | | | | | | | | |
| 4 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 274/2464 (11.1%) | 106/1736 (6.1%) | RR 1.28 (1.03 to 1.59) | 17 more per 1000 (from 2 more to 36 more) | HIGH | CRITICAL |
| | | | | | | | | 3.2% | | 8 more per 1,000 | | |
| | | | | | | | | 15.6% | | 43 more per 1,000 | | |

*Risk in the control group is reported as observed pooled, low and high-risks derived from included trials

Abbreviations: CI=confidence interval; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 44. KQ4 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | | Evidence Grade | Importance |
|--|------------------|------------------------|---------------|-------------------------|------------------------|----------------------|---------------------|-------------------|------------------------|---|------|----------------|------------|
| | | | | | | | No of patients | | Effect | | | | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Control | | Relative (95% CI) | Absolute | | | |
| Withdrawals due to ADR - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | | |
| 3 | randomised trial | no serious limitations | serious | no serious indirectness | serious | reporting bias | 732/5139 (14.2%) | 343/5096 (6.7%) | RR 2.30 (1.34 to 3.95) | 87 more per 1000 (from 23 more to 198 more) | LOW | IMPORTANT | |
| | | | | | | | | 1.0% | | 12 more per 1,000 | | | |
| | | | | | | | | 10.8% | | 140 more per 1,000 | | | |
| Withdrawals due to ADR - vs CCBs (follow-up 2-3 years) | | | | | | | | | | | | | |
| 2 | randomised trial | no serious limitations | serious | no serious indirectness | serious | reporting bias | 174/1495 (11.6%) | 128/1491 (8.6%) | RR 1.40 (0.92 to 2.12) | 34 more per 1000 (from 7 fewer to 96 more) | LOW | IMPORTANT | |
| | | | | | | | | 5.0% | | 19 more per 1,000 | | | |
| | | | | | | | | 13.1% | | 52 more per 1,000 | | | |
| Hypotension - IHD (follow-up 0.5-4.5 years) | | | | | | | | | | | | | |
| 3 | randomised trial | no serious limitations | serious | no serious indirectness | serious | reporting bias | 68/5490 (1.2%) | 26/5484 (0.5%) | RR 1.79 (0.68 to 4.71) | 5 fewer per 1000 (from 5 fewer to 5 fewer) | LOW | IMPORTANT | |
| | | | | | | | | 0.06% | | 0 fewer per 1,000 | | | |
| | | | | | | | | 3.2% | | 32 fewer per 1,000 | | | |
| Hypotension - vs CCBs (follow-up 2 years) | | | | | | | | | | | | | |
| 1 | randomised trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 64/673 (9.5%) | 22/663 (3.3%) | RR 2.87 (1.79 to 4.60) | 62 more per 1000 (from 26 more to 119 more) | HIGH | IMPORTANT | |
| Syncope - IHD (follow-up 4.5-4.8 years) | | | | | | | | | | | | | |
| 2 | randomised trial | no serious limitations | serious | no serious indirectness | serious | none | 203/8803 (2.3%) | 162/8784 (1.8%) | RR 1.24 (1.02 to 1.52) | 4 more per 1000 (from 0 more to 9 more) | LOW | IMPORTANT | |
| | | | | | | | | 0.2% | | 0 more per 1,000 | | | |
| | | | | | | | | 3.9% | | 9 more per 1,000 | | | |
| Cough - IHD (follow-up 2-4.8 years) | | | | | | | | | | | | | |
| 3 | randomised trial | no serious limitations | serious | no serious indirectness | serious | reporting bias | 1729/9476 (18.2%) | 1183/9439 (12.5%) | RR 1.67 (1.22 to 2.29) | 84 more per 1000 (from 28 more to 161 more) | LOW | IMPORTANT | |
| | | | | | | | | 0.2% | | 1 more per 1,000 | | | |
| | | | | | | | | 27.5% | | 184 more per 1,000 | | | |
| Cough - vs CCBs (follow-up 2 years) | | | | | | | | | | | | | |
| 1 | randomised trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 84/673 (12.5%) | 34/663 (5.1%) | RR 2.43 (1.66 to 3.57) | 73 more per 1000 (from 34 more to 131 more) | HIGH | IMPORTANT | |
| | | | | | | | | 0% | | 0 more per 1,000 | | | |

Appendix C: Additional Evidence Tables and Analyses

| Angioedema - IHD (follow-up 4.5-4.8 years) | | | | | | | | | | | | |
|--|------------------|------------------------|---------|-------------------------|---------|----------------|-----------------|-----------------|------------------------|---|-----|-----------|
| 2 | randomised trial | no serious limitations | serious | no serious indirectness | serious | none | 13/8803 (0.1%) | 6/8784 (0.1%) | RR 2.03 (0.75 to 5.47) | 1 more per 1000 (from 0 fewer to 4 more) | LOW | IMPORTANT |
| | | | | | | | | 0.2% | | 2 more per 1,000 | | |
| | | | | | | | | 1.2% | | 12 more per 1,000 | | |
| Hyperkalemia - IHD (follow-up 4.5-4.7 years) | | | | | | | | | | | | |
| 2 | randomised trial | no serious limitations | serious | no serious indirectness | serious | reporting bias | 506/7493 (6.8%) | 346/7544 (4.6%) | RR 1.71 (1.02 to 2.87) | 33 more per 1000 (from 1 more to 86 more) | LOW | IMPORTANT |
| | | | | | | | | 1.6% | | 11 more per 1,000 | | |
| | | | | | | | | 6.5% | | 46 more per 1,000 | | |

Abbreviations: CCB=calcium channel blocker; CI=confidence interval; CV=cardiovascular; IHD=ischemic heart disease; MI=myocardial infarction; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 45. KQ5 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | | Importance |
|--|------------------|------------------------|---------------|-------------------------|------------------------|----------------------|---------------------|-------------------|------------------------|--|----------------|------------|
| | | | | | | | No of patients | | Effect | | Evidence Grade | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Treatment | Control | Relative Risk (95% CI) | Absolute Risk | | |
| Study withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 2495/8502 (29.3%) | 2099/8576 (24.5%) | RR 1.20 (1.14 to 1.26) | 49 more per 1000 (from 34 more to 64 more) | HIGH | |
| Hypotension withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 406/8502 (4.8%) | 149/8576 (1.7%) | RR 2.75 (2.28 to 3.31) | 30 more per 1000 (from 22 more to 39 more) | HIGH | |
| Syncope withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | serious | none | 29/8502 (0.3%) | 15/8576 (0.2%) | RR 1.95 (1.06 to 3.60) | 2 more per 1000 (from 0 more to 5 more) | MODERATE | |
| Cough withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 392/8502 (4.6%) | 360/8576 (4.2%) | RR 1.10 (0.96 to 1.26) | 4 more per 1000 (from 2 fewer to 11 more) | HIGH | |
| Angioedema withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | very serious | none | 18/8502 (0.2%) | 25/8576 (0.3%) | RR 0.73 (0.40 to 1.32) | 1 fewer per 1000 (from 2 fewer to 1 more) | LOW | |
| Renal impairment withdrawals (follow-up 56 months) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single trial | no serious indirectness | no serious imprecision | none | 94/8502 (1.1%) | 60/8576 (0.7%) | RR 1.58 (1.15 to 2.18) | 4 more per 1000 (from 1 more to 8 more) | HIGH | |

Abbreviations: CI=confidence interval; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 46. KQ6 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | | | | Importance |
|---|------------------|------------------------|--------------------------|-------------------------|------------------------|----------------------|---------------------|-----------------|-------------------------|--|----------------|------------|
| | | | | | | | No of patients | | Effect | | Evidence Grade | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | Treatment | Control | Relative Risk (95% CI) | Absolute Risk | | |
| Study withdrawals (follow-up 0.5-3 years) | | | | | | | | | | | | |
| 3 | randomized trial | no serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | none | 230/1406 (16.4%) | 105/1397 (7.5%) | RR 2.17 (1.75 to 2.7) | 88 more per 1000 (from 56 more to 128 more) | HIGH | IMPORTANT |
| | | | | | | | | 0% | | 0 more per 1,000 | | |
| | | | | | | | | 8.1% | | 94 more per 1,000 | | |
| Hypotension (follow-up 3 years) | | | | | | | | | | | | |
| 1 | randomized trial | no serious limitations | single study | no serious indirectness | no serious imprecision | none | 154/120 (128.3%) | 70/1273 (5.5%) | RR 2.19 (1.67 to 2.87) | 65 more per 1000 (from 37 more to 103 more) | HIGH | IMPORTANT |
| Cough (follow-up 2.3-3 years) | | | | | | | | | | | | |
| 2 | randomized trial | no serious limitations | serious | no serious indirectness | very serious | none | 302/2158 (14%) | 143/2145 (6.7%) | RR 4.97 (0.58 to 42.95) | 266 more per 1000 (from 28 fewer to 1000 more) | LOW | IMPORTANT |
| | | | | | | | | 0.2% | | 7 more per 1,000 | | |
| | | | | | | | | 11.1% | | 440 more per 1,000 | | |

*Risk in the control group is reported as observed pooled, low and high-risks derived from included trials

Abbreviations: CI=confidence interval; RR=relative risk

Appendix C: Additional Evidence Tables and Analyses

Appendix Table 47. KQ7 - Strength of Evidence Grading

| Quality assessment | | | | | | | Summary of findings | | Importance |
|---|------------------|---------------------|--------------------------|-------------------------|------------------------|--|--|----------------|------------|
| | | | | | | | Findings | Evidence Grade | |
| No of studies | Design | Limitations | Inconsistency | Indirectness | Imprecision | Other considerations | | | |
| Sex impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 2 | randomized trial | serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the same in the two trials. | ACE inhibitors provide similar efficacy in males and females. | MODERATE | CRITICAL |
| Sex impact on benefits: ARB vs. placebo | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ARBs may not reduce the composite endpoint in females as much as males. | LOW | CRITICAL |
| Sex impact on benefits: ACE inhibitor vs. ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors may be superior to ARBs in females but similar in males. | LOW | CRITICAL |
| Sex impact on benefits: ACE inhibitor vs. ACE inhibitor + ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | Combination therapy may be superior to ACE inhibitors in females but similar in males. | LOW | CRITICAL |
| Sex impact on benefits: ACE inhibitor vs. CCB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors appear to be similar to CCBs in efficacy in either males or females. | LOW | CRITICAL |
| | | | | | | | | | |

Appendix C: Additional Evidence Tables and Analyses

| Age impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
|--|------------------|---------------------|--------------------------|-------------------------|------------------------|--|--|----------|----------|
| 2 | randomized trial | serious limitations | no serious imprecision | no serious indirectness | serious | Only the composite endpoint included in subgroup analyses. Composite endpoints not exactly the same in the two trials. Different age categories evaluated in different trials. | ACE inhibitors provide similar benefits in patients of different ages. | LOW | CRITICAL |
| Age impact on benefits: ARB vs. placebo | | | | | | | | | |
| 1 | randomized trial | serious limitations | single trial | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ARBs provide similar benefits in patients of different ages. | LOW | CRITICAL |
| Age impact on benefits: ACE inhibitor vs. ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single trial | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors provide similar benefits as ARBs in patients of different ages. | LOW | CRITICAL |
| Age impact on benefits: ACE inhibitor vs. ACE inhibitor + ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single trial | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors provide similar benefits as combination therapy in patients of different ages. | LOW | CRITICAL |
| Age impact on benefits: ACE inhibitor vs. CCB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single trial | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors provide similar benefits as calcium channel blockers in younger and older subjects. | LOW | CRITICAL |
| Diabetes mellitus impact on benefits: ACE inhibitor vs. Placebo | | | | | | | | | |
| 2 | randomized trial | serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses, which was not exactly the same in the two studies. Evaluated in subgroups (HOPE and EUROPA) and prespecified substudies (MICRO-HOPE, PERSUADE) from these trials. | ACE inhibitors provide similar benefits in those with and without diabetes mellitus. | MODERATE | CRITICAL |

Appendix C: Additional Evidence Tables and Analyses

| Diabetes mellitus impact on benefits: ARB vs. Placebo | | | | | | | | | |
|--|------------------|---------------------|--------------------------|-------------------------|------------------------|--|--|----------|----------|
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ARBs provide similar benefits in those with and without diabetes mellitus. | LOW | CRITICAL |
| Diabetes mellitus impact on benefits: ACE inhibitor vs. ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors provide similar benefits as ARBs in those with and without diabetes mellitus. | LOW | CRITICAL |
| Diabetes mellitus impact on benefits: ACE inhibitor vs. ACE inhibitor + ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER. | Combination therapy may be better than ACE inhibitors alone amongst those with diabetes mellitus but similar in non-diabetics. | LOW | CRITICAL |
| Diabetes mellitus impact on benefits: ACE inhibitor vs. CCB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses. Composite endpoint not exactly the one selected in the CER | ACE inhibitor therapy provides similar benefits as calcium channel blockers in subjects with diabetes. | LOW | CRITICAL |
| Renal dysfunction impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 3 | randomized trial | serious limitations | serious inconsistency | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the same in these two trials. The third trial evaluated total mortality instead of a composite endpoint. | ACE inhibitors may benefit those with renal dysfunction more than those without it. | LOW | CRITICAL |
| Hypertension impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 2 | randomized trial | serious limitations | no serious inconsistency | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoints not exactly the same in the two studies. | ACE inhibitors provide similar benefits to those with and without hypertension. | MODERATE | CRITICAL |

Appendix C: Additional Evidence Tables and Analyses

| Hypertension impact on benefits: ARB vs. placebo | | | | | | | | | |
|---|-------------------------------------|---------------------|--------------|-------------------------|------------------------|--|---|-----|----------|
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | ARBs provide similar benefits in those with and without hypertension. | LOW | CRITICAL |
| Hypertension impact on benefits: ACE inhibitor vs. ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | ACE inhibitors may provide more benefits to those with systolic hypertension while ARBs may provide more benefits to those with normal systolic blood pressure. | LOW | CRITICAL |
| Hypertension impact on benefits: ACE inhibitor vs. ACE inhibitor + ARB | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | Combination therapy with an ACE inhibitor and ARB may provide more benefits in lower and higher systolic blood pressure ranges but patients with modestly elevated systolic blood pressure may benefit more from ACE inhibitor therapy alone. | LOW | CRITICAL |
| Baseline risk impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 1 | Meta-analysis/ IPD meta-analysis | no limitations | single study | no serious indirectness | no serious imprecision | None | ACE inhibitors provide benefits in low, medium and high risk subjects. As baseline risk increases, the benefits derived from ACE inhibitor therapy might be accentuated. | LOW | CRITICAL |
| Baseline risk impact on benefits: ARB vs. placebo | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | ARBs may provide more benefits for those at lower baseline risk as compared to those at moderate to higher risk. | LOW | CRITICAL |
| Baseline risk impact on benefits: ACE inhibitors vs. ARBs | | | | | | | | | |
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | ACE inhibitor therapy might provide more benefits to those with moderate to high risk and ARBs may provide more benefits to those with lower baseline risk. | LOW | CRITICAL |

Appendix C: Additional Evidence Tables and Analyses

| Baseline risk impact on benefits: ACE inhibitor vs. ACE inhibitor + ARB | | | | | | | | | |
|---|---------------------------------|---------------------|--------------------------|-------------------------|------------------------|--|---|----------|----------|
| 1 | randomized trial | serious limitations | single study | no serious indirectness | no serious imprecision | Only the composite endpoint included in subgroup analyses in two trials. Composite endpoint not exactly the one selected in the CER. | Combination therapy with an ACE inhibitor + ARB provides similar benefits as an ACE inhibitor alone regardless of baseline risk. | LOW | CRITICAL |
| Antiplatelet therapy impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 1 | Meta-analysis/IPD meta-analysis | no limitations | no serious inconsistency | no serious indirectness | no serious imprecision | None | ACE inhibitors may provide more benefits to those without concurrent antiplatelet therapy as compared to those with antiplatelet therapy. ACE inhibitors provide significant benefits versus placebo in both subgroups. | MODERATE | CRITICAL |
| History of revascularization impact on benefits: ACE inhibitors vs. placebo | | | | | | | | | |
| 1 | Meta-analysis/IPD meta-analysis | no limitations | no serious inconsistency | no serious indirectness | no serious imprecision | None | ACE inhibitors may provide more benefits to those without a history of revascularization as compared to those with such a history. ACE inhibitors provide significant benefits versus placebo in both subgroups. | MODERATE | CRITICAL |
| Beta-blocker therapy impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 1 | Meta-analysis/IPD meta-analysis | No limitations | no serious inconsistency | no serious indirectness | no serious imprecision | None | ACE inhibitors provide similar benefits to those with and without beta-blocker therapy. ACE inhibitors provided significant benefits in those with and without beta-blockers. | MODERATE | CRITICAL |
| Lipid lowering therapy impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 1 | Meta-analysis/IPD meta-analysis | no limitations | no serious inconsistency | no serious indirectness | no serious imprecision | None | ACE inhibitors provide similar benefits to those with and lipid lowering therapy. ACE inhibitors provided significant benefits in those with and without beta-blockers. | MODERATE | CRITICAL |
| Vitamin E therapy impact on benefits: ACE inhibitor vs. placebo | | | | | | | | | |
| 1 | Randomized trial | no limitations | single study | no serious indirectness | serious imprecision | 95% confidence intervals and p-values not provided for this analysis. | ACE inhibitors provide similar benefits to those with and without vitamin E therapy. | LOW | CRITICAL |

Appendix D: Peer Reviewers

To Be Determined.

Appendix E: Abstraction Forms

Stable Ischemic Heart Disease Comparative Effectiveness Review

Study Characteristics

Ref ID:

| | | | |
|---------------|-------------------------------------|--------------------------|--|
| First Author: | | Citation: | |
| Language | Country(ies) Where Study Conducted: | Source of Study Funding: | |

Design Characteristics

| | | | | |
|--|---|------------------------------|--|------------------------------|
| Study Design: | | Randomization | Double-Blind? | Adequate? |
| <input type="checkbox"/> RCT - Parallel | <input type="checkbox"/> Obs - Registry | Adequate? | <input type="checkbox"/> Yes | <input type="checkbox"/> Yes |
| <input type="checkbox"/> RCT - Crossover | <input type="checkbox"/> Obs - Cohort | <input type="checkbox"/> Yes | <input type="checkbox"/> No | <input type="checkbox"/> No |
| <input type="checkbox"/> Obs - Case Control | <input type="checkbox"/> Other: | <input type="checkbox"/> No | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A |
| <input type="checkbox"/> Obs - Cross Sectional | | <input type="checkbox"/> N/A | | |
| Allocation Concealment? | Adequate? | Intention-to-Treat? | If no, What was other method? | |
| <input type="checkbox"/> Yes | <input type="checkbox"/> Yes | <input type="checkbox"/> Yes | <input type="checkbox"/> Prospective Study Design | |
| <input type="checkbox"/> No | <input type="checkbox"/> No | <input type="checkbox"/> No | <input type="checkbox"/> Propensity Score Matching | |
| <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> Propensity Score Adjusted | |
| | | | <input type="checkbox"/> Multivariate Analysis | |
| | | | <input type="checkbox"/> Other: | |

Study Population

| | | |
|------------------------------|-------------------------|------------------------------------|
| No. enrolled in Study: | No. Completed Study: | No. Withdrawals: |
| | | Intervention 1: |
| | | Reasons: |
| | | Intervention 2: |
| | | Reasons: |
| Run-in Period? | Describe Run-in Period: | % Removed During Run-in & Reasons: |
| <input type="checkbox"/> Yes | | |
| <input type="checkbox"/> No | | |
| <input type="checkbox"/> N/A | | |
| Intervention 1 (drug, dose): | | Intervention 2 (drug, dose): |
| Inclusion Criteria: | | |
| Exclusion Criteria: | | |
| Length of Study: | | Duration of Patient Follow-up: |

Appendix E: Abstraction Forms

Stable Ischemic Heart Disease Comparative Effectiveness Review

Baseline Characteristics

| | Intervention 1 | Intervention 2 | Total |
|--------------------------------------|----------------|----------------|-------|
| Males/Females: | | | |
| Average Age (years): | | | |
| White | | | |
| Hispanic | | | |
| African American | | | |
| Asian | | | |
| Other | | | |
| Average LVEF (%) | | | |
| Co-Morbidities | | | |
| Hx/o Angiographically Documented CAD | | | |
| Hx/o Myocardial Infarction | | | |
| Previous Revascularization | | | |
| CABG | | | |
| PTCA/PCI | | | |
| CABG or PTCA/PCI | | | |
| Hx/o Stable Angina | | | |
| Hx/o Unstable Angina | | | |
| Hx/o Stroke/TIA | | | |
| Hx/o Peripheral Vascular Disease | | | |
| Hx/o Diabetes | | | |
| Hx/o Renal Insufficiency | | | |
| Hx/o Hypertension | | | |
| Hx/o Left Ventricular Hypertrophy | | | |
| Hx/o Microalbuminuria | | | |
| Hx/o Smoking | | | |

Appendix E: Abstraction Forms

Stable Ischemic Heart Disease Comparative Effectiveness Review

Baseline Characteristics

| | Intervention 1 | Intervention 2 | Total |
|---------------------------------|----------------|----------------|-------|
| Systolic Blood Pressure (mmHg) | | | |
| Diastolic Blood Pressure (mmHg) | | | |
| Body Mass Index | | | |
| Total Cholesterol (mg/dl) | | | |
| LDL Cholesterol (mg/dL) | | | |
| HDL Cholesterol (mg/dL) | | | |
| Triglycerides (mg/dL) | | | |
| Glucose (mg/dl) | | | |
| Creatine (mg/dL) | | | |
| Potassium (mmol/L) | | | |
| Left Main | | | |
| Left Anterior Descending | | | |
| Left Circumflex | | | |
| Right Coronary Artery | | | |

Baseline Medical Therapies

| | | | |
|--------------------------|--|--|--|
| Beta-Blockers | | | |
| Calcium Channel Blockers | | | |
| Aspirin | | | |
| Clopidogrel/Ticlopidine | | | |
| Antiplatelet | | | |
| Diuretics | | | |
| Nitrates | | | |
| Statin | | | |
| Lipid-Lowering | | | |
| Digitalis | | | |

Appendix E: Abstraction Forms

Stable Ischemic Heart Disease Comparative Effectiveness Review

Efficacy Outcomes (Dichotomous)

| | Intervention 1 | | Intervention 2 | |
|--|----------------|------------------|----------------|------------------|
| | Number at risk | Number of events | Number at risk | Number of events |
| Primary Endpoint (list components) | | | | |
| Total Mortality | | | | |
| Cardiovascular Death | | | | |
| Total Myocardial Infarction | | | | |
| Fatal Myocardial Infarction | | | | |
| Non-Fatal Myocardial Infarction | | | | |
| Stroke | | | | |
| Composite (CV death, non-fatal MI, stroke) | | | | |
| Other Composite (list Components) | | | | |
| Other Composite (list Components) | | | | |
| Atrial Fibrillation | | | | |
| Hospitalization | | | | |
| No. of Ischemic Events | | | | |

Efficacy Outcomes (Continuous)

| | Intervention 1 | | Intervention 2 | |
|--------------------------------|----------------|-----------|----------------|-----------|
| | Baseline | Follow-up | Baseline | Follow-up |
| (n, mean \pm SD) | | | | |
| Exercise Time Before Ischemia | | | | |
| Quality of Life (Scale used:) | | | | |

Appendix E: Abstraction Forms

Stable Ischemic Heart Disease Comparative Effectiveness Review

Safety Outcomes (Dichotomous)

| | Intervention 1 | | Intervention 2 | |
|-----------------------------------|----------------|------------------|----------------|------------------|
| | Number at risk | Number of events | Number at risk | Number of events |
| Hyperkalemia | | | | |
| Cough | | | | |
| Angioedema | | | | |
| Syncope | | | | |
| Withdrawals Due to Adverse Events | | | | |
| Hypotension | | | | |
| Rash | | | | |
| Blood Dyscrasia's | | | | |